

#5245 Store at -20C

STAMBP Antibody



Cell Signaling
TECHNOLOGY®

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source:	UniProt ID:	Entrez-Gene Id:
WB, IP	H Mk	Endogenous	50	Rabbit	#O95630	10617

Product Usage Information

Application

Western Blotting
Immunoprecipitation

Dilution

1:1000
1:50

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at –20°C. Do not aliquot the antibody.

Specificity / Sensitivity

STAMBP Antibody detects endogenous levels of total STAMBP protein.

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Gly194 of human STAMBP protein. Antibodies are purified by protein A and peptide affinity chromatography.

Background

Protein ubiquitination and deubiquitination is a reversible process catalyzed by ubiquitinating enzymes (UBEs) and deubiquitinating enzymes (DUBs). Deubiquitinating enzymes (DUBs) function as ubiquitin-specific proteases and can be divided into five subfamilies based on catalytic domain structure. At least 14 members of the JAMM ubiquitin protease subfamily have been identified, including signal transducing adaptor molecule (STAM) binding protein (3). STAM-binding protein (STAMBP or AMSH) is an endosomal DUB that preferentially displays ubiquitin isopeptidase activity toward K63-linked chains, which is critically dependent upon its interaction with STAM (4,5). STAMBP interacts with the STAM adaptor protein and becomes integrated into the multivesicular body sorting machinery to help regulate endosomal trafficking and receptor tyrosine kinase stability by deubiquitinating target proteins (4,6). Evidence indicates that endosomal STAMBP antagonizes the ubiquitin-dependent trafficking of EGFR to the lysosomal compartment (7).

Background References

1. Nijman, S.M. et al. (2005) *Cell* 123, 773-86.
2. Nalepa, G. et al. (2006) *Nat Rev Drug Discov* 5, 596-613.
3. Kikuchi, K. et al. (2003) *Biochem Biophys Res Commun* 306, 637-43.
4. McCullough, J. et al. (2006) *Curr Biol* 16, 160-5.
5. Kim, M.S. et al. (2006) *Biochem Biophys Res Commun* 351, 612-8.
6. Agromayor, M. and Martin-Serrano, J. (2006) *J Biol Chem* 281, 23083-91.
7. McCullough, J. et al. (2004) *J Cell Biol* 166, 487-92.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting **IP:** Immunoprecipitation

Cross-Reactivity Key

H: human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster
X: Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse
GP: Guinea Pig **Rab:** rabbit **All:** all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.
All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.