5583 Store at -20C

Patents

DDR1 (D1G6) XP® Rabbit mAb



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: WB, IP, IHC-P, IF-IC	Reactivity: H M R Mk	Sensitivity: Endogenous	MW (kDa): 125	Source/Isotype: Rabbit IgG	UniProt ID: #Q08345	Entrez-Gene Id: 780	
Product Usage Information	Ар	plication			Dilution		
	We	stern Blotting			1:1000		
	Imr	munoprecipitation			1:100		
	Imr	munohistochemistry	(Paraffin)		1:3200 - 1:12800		
	Imr	munofluorescence (Immunocytochen	nistry)	1:1600 - 1:3200		
Storage	0.02	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody.					
Cuacificity / Canaiti		For a carrier free (BSA and azide free) version of this product see product #49876. DDR1 (D1G6) XP® Rabbit mAb detects endogeneous levels of total DDR1 protein.					
Specificity / Sensitivity		DDK1 (D100) AF Rabbit IIIAD detects endogeneous levels of total DDK1 protein.					
Source / Purificatio		Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Pro600 of human DDR1 protein.					
Background	theii DDF have abei	The discoidin domain receptors (DDRs) are receptor tyrosine kinases with a discoidin homology repeat in their extracellular domains, activated by binding to extracellular matrix collagens. So far, two mammalian DDRs have been identified: DDR1 and DDR2 (1). They are widely expressed in human tissues and may have roles in smooth muscle cell-mediated collagen remodeling (2). Research studies have implicated aberrant expression and signaling of DDRs in human diseases related to increased matrix degradation and remodeling, such as cardiovascular disease, liver fibrosis, and tumor invasion (1).					
Background Refere		 Vogel, W. (1999) FASEB J 13 Suppl, S77-82. Ferri, N. et al. (2004) Am J Pathol 164, 1575-85. 					

Species Reactivity Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS,

Applications Key WB: Western Blotting IP: Immunoprecipitation IHC-P: Immunohistochemistry (Paraffin)

IF-IC: Immunofluorescence (Immunocytochemistry)

Cross-Reactivity Key H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster

X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse

GP: Guinea Pig Rab: rabbit All: all species expected

Trademarks andCell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

XP is a registered trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more

information.

Limited Uses Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the

following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in

writing by a legally authorized representative of CST, are rejected and are of no force or effect.

DDR1 (D1G6) XP® Rabbit mAb (#5583) Datasheet Without Images Cell Signaling Technology

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.