

#5667 Store at -20C

NDRG2 Antibody


Cell Signaling
TECHNOLOGY®

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source:	UniProt ID:	Entrez-Gene Id:
WB	H M R	Endogenous	45	Rabbit	#Q9UN36	57447

Product Usage Information	Application Western Blotting	Dilution 1:1000
Storage	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at –20°C. Do not aliquot the antibody.	
Specificity / Sensitivity	NDRG2 Antibody recognizes endogenous levels of total NDRG2 protein.	
Source / Purification	Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Lys25 of human NDRG2 protein. Antibodies are purified by protein A and peptide affinity chromatography.	
Background	<p>The NDRG (N-Myc downstream-regulated gene) family consisting of NDRG1, NDRG2, NDRG3, and NDRG4 are structurally related proteins with roles in cell proliferation, differentiation, apoptosis, stress responses, and cell migration/metastasis (1-3). NDRG1 was originally identified as a protein that was upregulated in N-Myc knockout mice (1). Proteins in the NDRG family, particularly NDRG1 and NDRG2, have been reported to be down-regulated in various cancer tissues and have been suggested to function as a tumor suppressors (4,5).</p> <p>Expression of NDRG1 and NDRG2 is up-regulated by p53 and HIF-1, and contributes to apoptosis driven by those pathways (6-11). NDRG1 and NDRG2 are also associated with pathological conditions in the nervous system. Nonsense mutation of the NDRG1 gene has been shown to cause hereditary motor and sensory neuropathy-Lom (HMSNL), which is supported by studies demonstrating the role of NDRG1 in maintaining myelin sheaths and axonal survival (12, 13). Elevated expression of NDRG2 has been observed with Alzheimer's disease (14). Both NDRG1 and NDRG2 are phosphorylated at multiple sites by Akt and/or SGK1, although the precise physiological role of this phosphorylation is not known (15, 16).</p>	
Background References	1. Shimono, A. et al. (1999) <i>Mech Dev</i> 83, 39-52. 2. Qu, X. et al. (2002) <i>Mol Cell Biochem</i> 229, 35-44. 3. Zhou, R.H. et al. (2001) <i>Genomics</i> 73, 86-97. 4. Yao, L. et al. (2008) <i>Acta Biochim Biophys Sin (Shanghai)</i> 40, 625-35. 5. Ellen, T.P. et al. (2008) <i>Carcinogenesis</i> 29, 2-8. 6. Kurdiani, S.K. et al. (1998) <i>Cancer Res</i> 58, 4439-44. 7. Park, H. et al. (2000) <i>Biochem Biophys Res Commun</i> 276, 321-8. 8. Stein, S. et al. (2004) <i>J Biol Chem</i> 279, 48930-40. 9. Cangul, H. (2004) <i>BMC Genet</i> 5, 27. 10. Wang, L. et al. (2008) <i>Cell Physiol Biochem</i> 21, 239-50. 11. Liu, N. et al. (2008) <i>Nucleic Acids Res</i> 36, 5335-49. 12. Kalaydjieva, L. et al. (2000) <i>Am J Hum Genet</i> 67, 47-58. 13. Okuda, T. et al. (2004) <i>Mol Cell Biol</i> 24, 3949-56. 14. Mitchelmore, C. et al. (2004) <i>Neurobiol Dis</i> 16, 48-58. 15. Murray, J.T. et al. (2004) <i>Biochem J</i> 384, 477-88. 16. Burchfield, J.G. et al. (2004) <i>J Biol Chem</i> 279, 18623-32.	

Species Reactivity	Species reactivity is determined by testing in at least one approved application (e.g., western blot).
Western Blot Buffer	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.
Applications Key	WB: Western Blotting

Cross-Reactivity Key

H: human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster
X: Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse
GP: Guinea Pig **Rab:** rabbit **All:** all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.
All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.