

#5850 Store at -20C

Sall4 Antibody

Cell Signaling
TECHNOLOGY®Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source:	UniProt ID:	Entrez-Gene Id:
WB	H	Endogenous	80, 142	Rabbit	#Q9UJQ4	57167

Product Usage Information

Application

Western Blotting

Dilution

1:1000

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at –20°C. Do not aliquot the antibody.

Specificity / Sensitivity

Sall4 Antibody recognizes endogenous levels of total Sall4 protein, isoforms A and B. This antibody may also detect other Sall4 isoforms.

Species predicted to react based on 100% sequence homology:

Monkey

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues near the carboxy terminus of human Sall4 protein. Antibodies are purified by protein A and peptide affinity chromatography.

Background

Members of the *SALL* gene family encode putative zinc finger transcription factors highly expressed during development (1). Sall4 is expressed very early in development with other pluripotency regulators, such as Oct-4 and Nanog (2). Recent studies suggest Sall4 works as a master regulator that controls its own expression and the expression of Oct-4 in a transcriptional regulation feedback loop governing stem cell pluripotency and stem cell fate (2,3). Immunohistochemical studies indicate that Sall4 is a sensitive and specific diagnostic marker for primary germ cell tumors and yolk sac tumors (4,5). Research studies have shown that Sall4 is constitutively expressed in acute myeloid leukemia (AML) and is a probable effector of the Wnt/β-catenin signaling pathway in this disease (6). In addition, mutations in *SALL4* have been implicated in human malformation syndromes, including Duane-radial ray syndrome (Okihiro syndrome) and Acro-renal-ocular syndrome (7).

Background References

1. Sweetman, D. and Münsterberg, A. (2006) *Dev Biol* 293, 285-93.
2. Yang, J. et al. (2008) *Proc Natl Acad Sci USA* 105, 19756-61.
3. Yang, J. et al. (2010) *PLoS One* 5, e10766.
4. Mei, K. et al. (2009) *Mod Pathol* 22, 1628-36.
5. Miertus, J. et al. (2006) *Hum Genet* 119, 154-61.
6. Ma, Y. et al. (2006) *Blood* 108, 2726-35.
7. Al-Baradie, R. et al. (2002) *Am J Hum Genet* 71, 1195-9.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting

Cross-Reactivity Key

H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster
X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse
GP: Guinea Pig Rab: rabbit All: all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.