

#6404 Store at -20°C

## REPS1 (D6F4) Rabbit mAb



**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com

**Support:** 877-678-TECH (8324)

**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
WB, IP	H	Endogenous	125	Rabbit IgG	#Q96D71	85021

### Product Usage Information

#### Application

Western Blotting

Immunoprecipitation

#### Dilution

1:1000

1:200

### Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

### Specificity / Sensitivity

REPS1 (D6F4) Rabbit mAb recognizes endogenous levels of total REPS1 protein.

### Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Asn189 of human REPS1 protein.

### Background

REPS1 is a RalBP1-associated EH-homology domain containing protein. The sequence of REPS1 has an EH domain, followed by two proline-rich segments, and a C-terminal coiled-coil domain for binding to RalBP1 (1). The EH domain of REPS1 interacts with the NPF motif of Rab11-FIP2, mediates their colocalization to endosome vesicles, and influences EGFR endocytosis (2). The two proline-rich regions of REPS1 are important for binding to the SH3 domain of GRK/GRB2 and further regulate EGFR downstream signaling. The proline-rich regions of REPS1 have also been shown to interact with the SH3 domain of intersectin1 (ITSN1) and contribute to ITSN1/SGIP1/REPS1 complex formation on clathrin-coated pits (3). Three alternatively spliced isoforms of REPS1 have been identified.

### Background References

1. Yamaguchi, A. et al. (1997) *J Biol Chem* 272, 31230-4.
2. Cullis, D.N. et al. (2002) *J Biol Chem* 277, 49158-66.
3. Dergai, O. et al. (2010) *Biochem Biophys Res Commun* 402, 408-13.

### Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

### Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

### Applications Key

**WB:** Western Blotting **IP:** Immunoprecipitation

### Cross-Reactivity Key

**H:** human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster  
**X:** Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse  
**GP:** Guinea Pig **Rab:** rabbit **All:** all species expected

### Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.  
All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](http://cellsignal.com/trademarks) for more information.

### Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial

purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.