

SignalSilence® Stat3 siRNA I



10 µM in 300 µl
 (100 Transfections)

Orders ■ 877-616-CELL (2355)
 orders@cellsignal.com
Support ■ 877-678-TECH (8324)
 info@cellsignal.com
Web ■ www.cellsignal.com

rev. 02/11/16

For Research Use Only. Not For Use In Diagnostic Procedures.


Species Cross-Reactivity: H, M, R

Description: SignalSilence® Stat3 siRNA I from Cell Signaling Technology (CST) allows the researcher to specifically inhibit Stat3 expression using RNA interference, a method whereby gene expression can be selectively silenced through the delivery of double stranded RNA molecules into the cell. All SignalSilence® siRNA products from CST are rigorously tested in-house and have been shown to reduce target protein expression by western analysis.

Background: The Stat3 transcription factor is an important signaling molecule for many cytokines and growth-factor receptors (1) and is required for murine fetal development (2). Stat3 is constitutively activated in a number of human tumors (3,4) and possesses oncogenic potential (5) and anti-apoptotic activities (3). Stat3 is activated by phosphorylation at Tyr705, which induces dimerization, nuclear translocation and DNA binding (6,7). Transcriptional activation seems to be regulated by phosphorylation at Ser727 through the MAPK or mTOR pathways (8,9). Stat3 isoform expression appears to reflect biological function as the relative expression levels of Stat3α (86 kDa) and Stat3β (79 kDa) depend on cell type, ligand exposure or cell maturation stage (10). It is notable that Stat3β lacks the serine phosphorylation site within the carboxy-terminal transcriptional activation domain (8).

Directions for Use: CST recommends transfection with 100 nM Stat3 siRNA I 48 to 72 hours prior to cell lysis. For transfection procedure, follow protocol provided by the transfection reagent manufacturer. Please feel free to contact CST with any questions on use.

Quality Control: Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure appropriate coupling efficiency. The oligo is subsequently purified by affinity-solid phase extraction. The annealed RNA duplex is further analyzed by mass spectrometry to verify the exact composition of the duplex. Each lot is compared to the previous lot by mass spectrometry to ensure maximum lot-to-lot consistency.



Western blot analysis of extracts from HeLa cells, transfected with 100 nM SignalSilence® Control siRNA (Unconjugated) #6568 (-), SignalSilence® Stat3 siRNA I (+) or SignalSilence® Stat3 siRNA II #6582 (+), using Stat3 (79D7) Rabbit mAb #4904 and α-Tubulin (11H10) Rabbit mAb #2125. The Stat3 (79D7) Rabbit mAb confirms silencing of Stat3 expression, while the α-Tubulin (11H10) Rabbit mAb is used to control for loading and specificity of Stat3 siRNA.

Entrez-Gene ID #6774
Swiss-Prot Acc. #P40763

Storage: Stat3 siRNA I is supplied in RNase-free water. Aliquot and store at -20°C.

Please visit www.cellsignal.com for a complete listing of recommended companion products.

Background References:

- (1) Heim, M.H. (1999) *J. Recept. Signal Transduct. Res.* 19, 75–120.
- (2) Takeda, K. et al. (1997) *Proc. Natl. Acad. Sci. USA* 94, 3801–3804.
- (3) Cattlett-Falcone, R. et al. (1999) *Immunity* 10, 105–115.
- (4) Garcia, R. and Jove, R. (1998) *J. Biomed. Sci.* 5, 79–85.
- (5) Bromberg, J.F. et al. (1999) *Cell* 98, 295–303.
- (6) Darnell Jr., J.E. et al. (1994) *Science* 264, 1415–1421.
- (7) Ihle, J.N. (1995) *Nature* 377, 591–594.
- (8) Wen, Z. et al. (1995) *Cell* 82, 241–250.
- (9) Yokogami, K. et al. (2000) *Curr. Biol.* 10, 47–50.
- (10) Biethahn, S. et al. (1999) *Exp. Hematol.* 27, 885–894.