

#6954 Store at -20C

CdGAP Antibody

**Cell Signaling**
TECHNOLOGY®**Orders:** 877-616-CELL (2355)
orders@cellsignal.com**Support:** 877-678-TECH (8324)**Web:** info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source:	UniProt ID:	Entrez-Gene Id:
WB	H M	Endogenous	250	Rabbit	#Q2M1Z3	57514

Product Usage Information**Application**

Western Blotting

Dilution

1:1000

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at –20°C. Do not aliquot the antibody.

Specificity / Sensitivity

CdGAP Antibody detects endogenous levels of total CdGAP protein. In certain cell lines, CdGAP Antibody recognizes a 125 kDa band of unknown origin.

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues near the central region of human CdGAP protein. Antibodies are purified by protein A and peptide affinity chromatography.

Background

The Rho family of small GTPases, including Rho, Rac, and Cdc42, act as molecular switches that regulate processes such as cell migration, adhesion, proliferation, and differentiation. They are activated by guanine nucleotide exchange factors (GEFs), which catalyze the exchange of bound GDP for GTP, and inhibited by GTPase activating proteins (GAPs), which catalyze the hydrolysis of GTP to GDP (1). The serine- and proline-rich GAP protein, Cdc42 GAP (CdGAP), has been shown to be a negative regulator of both Cdc42 and Rac1, but not RhoA (2,3). This protein contains three domains: an amino-terminal GAP domain, a central domain, and a carboxy-terminal proline-rich domain containing five Src homology 3 (SH3)-binding sites. It is suggested that threonine and serine phosphorylation within the proline-rich domain likely alters protein-protein interactions and determines the localization of CdGAP (4). Phosphorylation of CdGAP on threonine 776 by both ERK-1 and GSK-3 has been shown to negatively regulate protein activity, possibly by inducing a conformational change within the protein disrupting its ability to bind SH3 domains (4,5). Upregulation of CdGAP has been shown to increase cell proliferation and it has been suggested that this protein may play a role in TGF-β-induced cell growth, motility, and invasion in some breast cancer cells (6).

Background References

1. Takai, Y. et al. (2001) *Physiol Rev* 81, 153-208.
2. Tcherkezian, J. et al. (2006) *Biol Cell* 98, 445-56.
3. Lamarche-Vane, N. and Hall, A. (1998) *J Biol Chem* 273, 29172-7.
4. Tcherkezian, J. et al. (2005) *Mol Cell Biol* 25, 6314-29.
5. Danek, E.I. et al. (2007) *J Biol Chem* 282, 3624-31.
6. He, Y. et al. (2011) *Oncogene* 30, 1032-45.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key**WB:** Western Blotting**Cross-Reactivity Key**

H: human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster
X: Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse
GP: Guinea Pig **Rab:** rabbit **All:** all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.**Limited Uses**

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.