Phospho-Jak2 (Tyr1008) (D4A8) Rabbit mAb



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

| Applications: WB | Reactivity: H M | Sensitivity: Endogenous | MW (kDa): 125 | Source/Isotype: Rabbit IgG | UniProt ID: #O60674 | Entrez-Gene Id 3717 | |
|---|--------------------|---|---|-------------------------------|------------------------|------------------------|--|
| Product Usage Information | Ар | Application | | | Dilution | | |
| | We | Western Blotting | | | 1:1000 | | |
| Storage | | Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μ g/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20° C. Do not aliquot the antibody. | | | | | |
| phosphorylated at Tyr1008. T | | 008. This antibod | 4A8) Rabbit mAb recognizes endogenous levels of Jak2 protein only when This antibody also reacts with Jak2 when dually phosphorylated at Tyr1007 ty was not observed with other Jak family members by western blot. | | | | |
| Species predicted to react based on 100% sequence homology: | | | Bovine, Pig | | | | |

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Tyr1008 of human Jak2 protein.

Background

Members of the Janus family of tyrosine kinases (Jak1, Jak2, Jak3, and Tyk2) are activated by ligands binding to a number of associated cytokine receptors (1). Upon cytokine receptor activation, Jak proteins become autophosphorylated and phosphorylate their associated receptors to provide multiple binding sites for signaling proteins. These associated signaling proteins, such as Stats (2), Shc (3), insulin receptor substrates (4), and focal adhesion kinase (FAK) (5), typically contain SH2 or other phospho-tyrosine-binding domains.

Jak2 signaling is associated with a number of cytokines, growth factors, and hormones including IL-3, IL-5, IL-6, granulocyte-macrophage colony-stimulating factor (GM-CSF), erythropoietin (EPO), thrombopoietin (TPO) growth hormone, prolactin, and leptin (6-13). The oncogenic potential of Jak2 has been realized though translocations and point mutations resulting in its enhanced, de-regualated kinase activity, making Jak2 a potential therapeutic target. Jak2 gene translocations resulting in fusions with the TEL (TEL-JAK2) and PCM1 (PCM1-JAK2) have been found in leukemia patients (14,15). An activating point mutation in Jak2 resulting in a valine to phenylalanine switch at position 617 (V617F) has been implicated in myeloproliferative disorders including polycythemia vera and essential thrombocythemia (16).

Background References

- 1. Leonard, W.J. and O'Shea, J.J. (1998) Annu Rev Immunol 16, 293-322.
- 2. Darnell, J.E. (1997) Science 277, 1630-5.
- 3. VanderKuur, J. et al. (1995) *J Biol Chem* 270, 7587-93.
- 4. Argetsinger, L.S. et al. (1995) J Biol Chem 270, 14685-92.
- 5. Zhu, T. et al. (1998) J Biol Chem 273, 10682-9.
- 6. Quelle, F.W. et al. (1994) Mol Cell Biol 14, 4335-41.
- 7. Witthuhn, B.A. et al. (1993) Cell 74, 227-36.
- 8. Tortolani, P.J. et al. (1995) Blood 85, 3444-51.
- 9. Pallard, C. et al. (1995) EMBO J 14, 2847-56.
- 10. Narazaki, M. et al. (1994) Proc Natl Acad Sci USA 91, 2285-9.
- 11. Argetsinger, L.S. et al. (1993) Cell 74, 237-44.
- 12. Rui, H. et al. (1994) J Biol Chem 269, 5364-8.
- 13. Ghilardi, N. and Skoda, R.C. (1997) Mol Endocrinol 11, 393-9.
- 14. Lacronique, V. et al. (1997) Science 278, 1309-12.
- 15. Reiter, A. et al. (2005) Cancer Res 65, 2662-7.
- 16. Kralovics, R. et al. (2005) N Engl J Med 352, 1779-90.

3/23/24, 11:25 AM Phospho-Jak2 (Tyr1008) (D4A8) Rabbit mAb (#8082) Datasheet Without Images Cell Signaling Technology

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting

Cross-Reactivity Key

 $\textbf{H:} \ \text{human} \ \textbf{M:} \ \text{mouse} \ \textbf{R:} \ \text{rat} \ \textbf{Hm:} \ \text{hamster} \ \textbf{Mk:} \ \text{monkey} \ \textbf{Vir:} \ \text{virus} \ \textbf{Mi:} \ \text{mink} \ \textbf{C:} \ \text{chicken} \ \textbf{Dm:} \ \textbf{D.} \ \text{melanogaster}$

X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse

GP: Guinea Pig Rab: rabbit All: all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

Jak antibodies produced under license (granting certain rights including those under U.S. Patent No.

5,658,791) from Chemicon International, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more

information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.