

#8097 Store at -20°C

TECPR1 (D6C10) Rabbit mAb



Cell Signaling
TECHNOLOGY®

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
WB	H M R	Endogenous	130	Rabbit IgG	#Q7Z6L1	25851

Product Usage Information	Application Western Blotting	Dilution 1:1000
Storage	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.	
Specificity / Sensitivity	TECPR1 (D6C10) Rabbit mAb recognizes endogenous levels of total TECPR1 protein. This antibody does not cross-react with TECPR2. A background band of unknown origin around 200 kDa is detected in some cell lines.	
Source / Purification	Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding His900 of human TECPR1 protein.	
Background	Tectonin β-propeller repeat protein (TECPR1) was first identified as DKFZP434B0335, a human ortholog of the yeast protein Pex23p (1). TECPR1 contains a pleckstrin homology (PH) domain, β-propeller domain, and dysferlin domains. Research studies have shown that elevated expression of TECPR1 may be a potential marker for prostate cancer (2). In several independent studies, TECPR1 was shown to play a role in autophagy through interaction with Atg5 (3-5). Atg5 is a protein that is conjugated to the ubiquitin-like protein Atg12 and plays an essential role in autophagy (6). Initial studies suggested that TECPR1 plays a role in selective autophagy processes by targeting bacterial pathogens, as well as damaged mitochondria and protein aggregates (4). TECPR1 appears to be important for autophagosome maturation and promotes autophagosome fusion with the lysosome (5). Deletion of TECPR1 leads to an accumulation of autophagosomes (5).	
Background References	<ol style="list-style-type: none"> 1. Jeynov, B. et al. (2006) <i>FEBS Lett</i> 580, 5917-24. 2. Bull, J.H. et al. (2001) <i>Br J Cancer</i> 84, 1512-9. 3. Behrends, C. et al. (2010) <i>Nature</i> 466, 68-76. 4. Ogawa, M. et al. (2011) <i>Cell Host Microbe</i> 9, 376-89. 5. Chen, D. et al. (2012) <i>Mol Cell</i> 45, 629-41. 6. Mizushima, N. et al. (1998) <i>Nature</i> 395, 395-8. 	

Species Reactivity	Species reactivity is determined by testing in at least one approved application (e.g., western blot).
Western Blot Buffer	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.
Applications Key	WB: Western Blotting
Cross-Reactivity Key	H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse GP: Guinea Pig Rab: rabbit All: all species expected
Trademarks and Patents	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.
Limited Uses	Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.