Phospho-Stat3 (Tyr705) (D3A7) XP[®] Rabbit mAb (PE Conjugate)



Orders: 877-616-CELL (2355)

orders@cellsignal.com

877-678-TECH (8324) Support:

Web: info@cellsignal.com

cellsignal.com

3 Track Lane | Danvers | Massachusetts | 01923 | LISA

		3 Trask Lane Danvers Massachusetts 01923 U				
For Research Use Only. Not for Use in Diagnostic Procedures.						
Applications: FC-FP	Reactivity: H M R Mk	Sensitivity: Endogenous	Source/Isotype: Rabbit IgG	UniProt ID: #P40763	Entrez-Gene Id: 6774	
Product Usage	Ар	plication		Dilution		
Information	Flo	w Cytometry (Fixe	ed/Permeabilized)		1:50	
Storage	•	Supplied in PBS (pH 7.2), less than 0.1% sodium azide and 2 mg/ml BSA. Store at 4°C. Do not aliquot the antibodies. Protect from light. Do not freeze.				
Specificity / Sens	whe	Phospho-Stat3 (Tyr705) (D3A7) XP [®] Rabbit mAb (PE Conjugate) detects endogenous levels of Stat3 only when phosphorylated at Tyr705. This antibody does not cross-react with phospho-EGFR or the corresponding phospho-tyrosines of other Stat proteins.				
Species predicted react based on 10 sequence homological contracts and contracts are contracted by the contract of the contrac	00%	nster, Bovine, Pig,	Horse			
Source / Purificat		Monoclonal antibody is produced by immunizing animals with a synthetic phosphopeptide corresponding to residues surrounding Tyr705 of mouse Stat3 protein.				
Product Descripti	flow	This Cell Signaling Technology antibody is conjugated to phycoerythrin (PE) and tested in-house for direct flow cytometry analysis in human cells. The antibody is expected to exhibit the same species cross-reactivity as the unconjugated Phospho-Stat3 (Tyr705) (D3A7) XP® Rabbit mAb #9145.				

Background

The Stat3 transcription factor is an important signaling molecule for many cytokines and growth factor receptors (1) and is required for murine fetal development (2). Research studies have shown that Stat3 is constitutively activated in a number of human tumors (3,4) and possesses oncogenic potential (5) and antiapoptotic activities (3). Stat3 is activated by phosphorylation at Tyr705, which induces dimerization, nuclear translocation, and DNA binding (6,7). Transcriptional activation seems to be regulated by phosphorylation at Ser727 through the MAPK or mTOR pathways (8,9). Stat3 isoform expression appears to reflect biological function as the relative expression levels of Stat3α (86 kDa) and Stat3β (79 kDa) depend on cell type, ligand exposure, or cell maturation stage (10). It is notable that Stat3β lacks the serine phosphorylation site within the carboxy-terminal transcriptional activation domain (8).

Background References

- 1. Heim, M.H. (2001) J Recept Signal Transduct Res 19, 75-120.
- 2. Takeda, K. et al. (1997) Proc Natl Acad Sci U S A 94, 3801-4.
- 3. Catlett-Falcone, R. et al. (1999) Immunity 10, 105-15.
- 4. Garcia, R. and Jove, R. (1998) J Biomed Sci 5, 79-85.
- 5. Bromberg, J.F. et al. (1999) Cell 98, 295-303.
- 6. Darnell, J.E. et al. (1994) Science 264, 1415-21.
- 7. Ihle, J.N. (1995) Nature 377, 591-4.
- 8. Wen, Z. et al. (1995) Cell 82, 241-50.
- 9. Yokogami, K. et al. (2000) Curr Biol 10, 47-50.

10. Biethahn, S. et al. (1999) Exp Hematol 27, 885-94.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Applications Key

Cross-Reactivity Key

FC-FP: Flow Cytometry (Fixed/Permeabilized)

H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse GP: Guinea Pig Rab: rabbit All: all species expected

Trademarks and Patents

Limited Uses

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

XP is a registered trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.