

#8204 Store at -20°C

PhosphoPlus® Stat3 (Tyr705) Antibody Duet



Cell Signaling
TECHNOLOGY®

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

UniProt ID:
#P40763

Entrez-Gene Id:
6774

Product Includes	Product #	Quantity	Mol. Wt.	Isotype/Source
Phospho-Stat3 (Tyr705) (D3A7) XP® Rabbit mAb	9145	100 µl	79, 86 kDa	Rabbit IgG
Stat3 (D3Z2G) Rabbit mAb	12640	100 µl	79, 86 kDa	Rabbit IgG

Please visit [cellsignal.com](https://www.cellsignal.com) for individual component applications, species cross-reactivity, dilutions, protocols, and additional product information.

Description

PhosphoPlus® Duets from Cell Signaling Technology (CST) provide a means to assess protein activation status. Each Duet contains an activation-state and total protein antibody to your target of interest. These antibodies have been selected from CST's product offering based upon superior performance in specified applications.

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. *Do not aliquot the antibody.*

Background

The Stat3 transcription factor is an important signaling molecule for many cytokines and growth factor receptors (1) and is required for murine fetal development (2). Research studies have shown that Stat3 is constitutively activated in a number of human tumors (3,4) and possesses oncogenic potential (5) and anti-apoptotic activities (3). Stat3 is activated by phosphorylation at Tyr705, which induces dimerization, nuclear translocation, and DNA binding (6,7). Transcriptional activation seems to be regulated by phosphorylation at Ser727 through the MAPK or mTOR pathways (8,9). Stat3 isoform expression appears to reflect biological function as the relative expression levels of Stat3α (86 kDa) and Stat3β (79 kDa) depend on cell type, ligand exposure, or cell maturation stage (10). It is notable that Stat3β lacks the serine phosphorylation site within the carboxy-terminal transcriptional activation domain (8).

Background References

1. Heim, M.H. (2001) *J Recept Signal Transduct Res* 19, 75-120.
2. Takeda, K. et al. (1997) *Proc Natl Acad Sci U S A* 94, 3801-4.
3. Catlett-Falcone, R. et al. (1999) *Immunity* 10, 105-15.
4. Garcia, R. and Jove, R. (1998) *J Biomed Sci* 5, 79-85.
5. Bromberg, J.F. et al. (1999) *Cell* 98, 295-303.
6. Darnell, J.E. et al. (1994) *Science* 264, 1415-21.
7. Ihle, J.N. (1995) *Nature* 377, 591-4.
8. Wen, Z. et al. (1995) *Cell* 82, 241-50.
9. Yokogami, K. et al. (2000) *Curr Biol* 10, 47-50.
10. Biethahn, S. et al. (1999) *Exp Hematol* 27, 885-94.

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.
PhosphoPlus is a registered trademark of Cell Signaling Technology, Inc.
U.S. Patent No. 7,429,487, foreign equivalents, and child patents deriving therefrom.
All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](https://www.cellsignal.com/trademarks) for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the

end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.