

#8225 Store at -20°C

CACYBP (D43G11) Rabbit mAb**Cell Signaling**
TECHNOLOGY®**Orders:** 877-616-CELL (2355)
orders@cellsignal.com**Support:** 877-678-TECH (8324)**Web:** info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
WB, IHC-P	H M R Mk	Endogenous	26	Rabbit IgG	#Q9HB71	27101

Product Usage Information**Application**Western Blotting
Immunohistochemistry (Paraffin)**Dilution**1:1000
1:50**Storage**

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Specificity / Sensitivity

CACYBP (D43G11) Rabbit mAb recognizes endogenous levels of total CACYBP protein.

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Arg162 of human CACYBP protein.

Background

Calcyclin-binding protein or Siah-1-interacting protein (CACYBP/SIP) is a component of the ubiquitin E3 ligase complex that also contains Siah1, Skp1, and Ebi (1). CACYBP regulates β -catenin turnover and plays an important role in thymocyte development (2). CACYBP also binds to tubulin and may be involved in cytoskeletal regulation (3,4). It is highly expressed in neurons, and its cellular localization may be regulated by Ca^{2+} (5,6). Retinoic acid treatment of the neuroblastoma cell line SH-SY5Y induces translocation of CACYBP to the nucleus and seems to be correlated with phosphorylation of CACYBP on serine residues (7). Recent studies also suggest that CACYBP may possess phosphatase activity (8), and that it can bind and dephosphorylate Erk1/2 (8,9).

Background References

1. Matsuzawa, S.I. and Reed, J.C. (2001) *Mol Cell* 7, 915-26.
2. Fukushima, T. et al. (2006) *Immunity* 24, 29-39.
3. Schneider, G. et al. (2007) *Biochim Biophys Acta* 1773, 1628-36.
4. Filipek, A. et al. (2008) *J Neural Transm* 115, 1257-64.
5. Jastrzebska, B. et al. (2000) *J Histochem Cytochem* 48, 1195-202.
6. Filipek, A. et al. (2002) *J Biol Chem* 277, 21103-9.
7. Wu, J. et al. (2003) *J Biochem Mol Biol* 36, 354-8.
8. Kilanczyk, E. et al. (2011) *Biochem Biophys Res Commun* 404, 179-83.
9. Kilanczyk, E. et al. (2009) *Biochem Biophys Res Commun* 380, 54-9.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key**WB:** Western Blotting **IHC-P:** Immunohistochemistry (Paraffin)**Cross-Reactivity Key**

H: human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster
X: Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse
GP: Guinea Pig **Rab:** rabbit **All:** all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.
 All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.