#8457 Store at -20C

β-Actin (D6A8) Rabbit mAb



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: WB, IF-IC, FC-FP	Reactivity: H M R Mk Dm Z	Sensitivity: Endogenous	MW (kDa): 45	Source/Isotype: Rabbit IgG	UniProt ID: #P60709	Entrez-Gene Id: 60
Product Usage Information	Application				Dilution	
	Western Blotting				1:1000	
	Immunofluorescence (Immunocytochemistry)				1:100 - 1:200	
	Flow Cytometry (Fixed/Permeabilized)				1:50 - 1:200	
Storage	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μ g/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody.					
	For a carrier free (BSA and azide free) version of this product see product #19069.					

Specificity / Sensitivity

 β -Actin (D6A8) Rabbit mAb recognizes endogenous levels of total β -actin protein. Due to the high sequence identity between the cytoplasmic actin isoforms, β -actin and cytoplasmic γ -actin, this antibody may cross-react with cytoplasmic γ -actin. It does not cross-react with α -skeletal, α -cardiac, α -vascular smooth, or γ -enteric smooth muscle isoforms.

Species predicted to react based on 100% sequence homology:

Hamster, Chicken, Bovine, Pig

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues near the amino terminus of human β -actin protein.

Background

Actin, a ubiquitous eukaryotic protein, is the major component of the cytoskeleton. At least six isoforms are known in mammals. Nonmuscle β - and γ -actin, also known as cytoplasmic actin, are ubiquitously expressed, controlling cell structure and motility (1). While all actin isoforms are highly homologous, cytoplasmic β - and γ -actin protein sequences differ by only four biochemically similar amino acids (2). For this reason, antibodies raised to β -actin may cross-react with γ -actin, and vice versa. α -cardiac and α -skeletal actin are expressed in striated cardiac and skeletal muscles, respectively; two smooth muscle actins, α - and γ -actin, are found primarily in vascular smooth muscle and enteric smooth muscle, respectively. These actin isoforms regulate the contractile potential of muscle cells (1). Actin exists mainly as a fibrous polymer, F-actin. In response to cytoskeletal reorganizing signals during processes such as cytokinesis, endocytosis, or stress, cofilin promotes fragmentation and depolymerization of F-actin, resulting in an increase in the monomeric globular form, G-actin (3). The ARP2/3 complex stabilizes F-actin fragments and promotes formation of new actin filaments (3). Research studies have shown that actin is hyperphosphorylated in primary breast tumors (4). Cleavage of actin under apoptotic conditions has been observed *in vitro* and in cardiac and skeletal muscle, as shown in research studies (5-7). Actin cleavage by caspase-3 may accelerate ubiquitin/proteasome-dependent muscle proteolysis (7).

Background References

- 1. Herman, I.M. (1993) Curr. Opin. Cell Biol. 5, 48-55.
- 2. Perrin, B.J. and Ervasti, J.M. (2010) Cytoskeleton (Hoboken) 67, 630-4.
- 3. Condeelis, J. (2001) Trends Cell Biol 11, 288-93.
- 4. Lim, Y.P. et al. (2004) Clin Cancer Res 10, 3980-7.
- 5. Kayalar, C. et al. (1996) *Proc Natl Acad Sci U S A* 93, 2234-8.
- 6. Communal, C. et al. (2002) Proc Natl Acad Sci U S A 99, 6252-6.
- 7. Du, J. et al. (2004) J Clin Invest 113, 115-23.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

1/1/24. 3:03 PM

Western Blot Buffer

β-Actin (D6A8) Rabbit mAb (#8457) Datasheet Without Images Cell Signaling Technology

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting IF-IC: Immunofluorescence (Immunocytochemistry)

FC-FP: Flow Cytometry (Fixed/Permeabilized)

Cross-Reactivity Key

H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster

X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse

GP: Guinea Pig Rab: rabbit All: all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. Alexa Fluor is a registered trademark of Life Technologies Corporation.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more

information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.