

#8532 Store at -20°C

FYVE-CENT Antibody


Cell Signaling
TECHNOLOGY®

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source:	UniProt ID:	Entrez-Gene Id:
WB, IP	H	Endogenous	285	Rabbit	#Q68DK2	23503

Product Usage Information

Application

Western Blotting

Immunoprecipitation

Dilution

1:1000

1:50

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at –20°C. Do not aliquot the antibody.

Specificity / Sensitivity

FYVE-CENT Antibody recognizes endogenous levels of total FYVE-CENT protein.

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Leu595 of human FYVE-CENT protein. Antibodies are purified by protein A and peptide affinity chromatography.

Background

FYVE-CENT (ZFYE26) contains an FYVE zinc finger domain characterized by conserved R(R/RK) and HHCRxCG motifs (1). Phosphatidylinositol 3 phosphate binds to FYVE-CENT via the FYVE domain and this binding is crucial in phosphoinositide (PtdIns3P)-regulated cytokinesis (1-3). Research evidence suggests that during cytokinesis, FYVE-CENT directly interacts with TTC19 and is recruited to the midbody by the kinesin protein KIF13A (4). FYVE-CENT may also be involved in the tumor suppressor function of Beclin-1 (5). Researchers have linked the defects in FYVE-CENT to hereditary spastic paraplegia, a condition characterized by deterioration of the corpus callosum of the brain (6).

Background References

1. Stenmark, H. and Aasland, R. (1999) *J Cell Sci* 112 (Pt 23), 4175-83.
2. Stenmark, H. et al. (2002) *FEBS Lett* 513, 77-84.
3. Gaullier, J.M. et al. (1999) *Chem Phys Lipids* 98, 87-94.
4. Sagona, A.P. et al. (2011) *PLoS One* 6, e17086.
5. Sagona, A.P. et al. (2010) *Nat Cell Biol* 12, 362-71.
6. Denora, P.S. et al. (2009) *J Neurol Sci* 277, 22-5.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting **IP:** Immunoprecipitation

Cross-Reactivity Key

H: human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster **X:** Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse **GP:** Guinea Pig **Rab:** rabbit **All:** all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner

that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.