Store at -200

HER2/ErbB2 (D8F12) XP® Rabbit mAb (Biotinylated)



Orders:

877-616-CELL (2355) orders@cellsignal.com

Support:

877-678-TECH (8324)

Web:

info@cellsignal.com cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures. Dogotivity

Applications: WB	Reactivity: H M	Sensitivity: Endogenous	MW (kDa): 185	Source/Isotype: Rabbit IgG	UniProt ID: #P04626	Entrez-Gene Id: 2064
Product Usage	Ар	plication			Dilution	
Information	We	stern Blotting			1:1000	
Storage		Supplied in 136 mM NaCl, 2.6 mM KCl, 12 mM sodium phosphate (pH 7.4) dibasic, 2 mg/ml BSA, and 50% glycerol. Store at -20° C. Do not aliquot the antibodies.				
Specificity / Sen		HER2/ErbB2 (D8F12) ${\sf XP}^{\it \&}$ Rabbit mAb (Biotinylated) detects endogenous levels of total HER2/ErbB2 protein.				
Species predicte react based on 1 sequence homo	.00%					
Source / Purification		Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues near the amino terminus of human HER2/ErbB2 protein.				
Product Descrip	antil	nis Cell Signaling Technology antibody is conjugated to biotin under optimal conditions. The biotinylated ntibody is expected to exhibit the same species cross-reactivity as the unconjugated HER2/ErbB2 NRF12) XP® Rabbit mAb #4290.				

Background

MW (kDa)

The ErbB2 (HER2) proto-oncogene encodes a 185 kDa transmembrane, receptor-like glycoprotein with intrinsic tyrosine kinase activity (1). While ErbB2 lacks an identified ligand, ErbB2 kinase activity can be activated in the absence of a ligand when overexpressed and through heteromeric associations with other ErbB family members (2). Amplification of the ErbB2 gene and overexpression of its product are detected in almost 40% of human breast cancers (3). Binding of the c-Cbl ubiquitin ligase to ErbB2 at Tyr1112 leads to ErbB2 poly-ubiquitination and enhances degradation of this kinase (4). ErbB2 is a key therapeutic target in the treatment of breast cancer and other carcinomas and targeting the regulation of ErbB2 degradation by the c-Cbl-regulated proteolytic pathway is one potential therapeutic strategy. Phosphorylation of the kinase domain residue Tyr877 of ErbB2 (homologous to Tyr416 of pp60c-Src) may be involved in regulating ErbB2 biological activity. The major autophosphorylation sites in ErbB2 are Tyr1248 and

185

transduction pathway (1,5).

Background References

- 1. Muthuswamy, S.K. et al. (1999) Mol Cell Biol 19, 6845-57.
- 2. Qian, X. et al. (1994) Proc Natl Acad Sci USA 91, 1500-4.
- 3. Dittadi, R. and Gion, M. (2000) J Natl Cancer Inst 92, 1443-4.
- 4. Klapper, L.N. et al. (2000) Cancer Res 60, 3384-8.
- 5. Kwon, Y.K. et al. (1997) J Neurosci 17, 8293-9.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Tyr1221/1222; phosphorylation of these sites couples ErbB2 to the Ras-Raf-MAP kinase signal

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting

1/1/24, 8:27 AM

HER2/ErbB2 (D8F12) XP® Rabbit mAb (Biotinylated) (#8609) Datasheet Without Images Cell Signaling Tech...

Cross-Reactivity Key

H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster X: Xenopus Z: zebrafish B: bovine Dq: dog Pq: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse

GP: Guinea Pig Rab: rabbit All: all species expected

Trademarks and Patents

Limited Uses

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

XP is a registered trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.