

#8658 Store at -20°C

Notch Receptor Interaction Antibody Sampler Kit

1 Kit (9 x 20 microliters)



Cell Signaling
TECHNOLOGY®

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Product Includes	Product #	Quantity	Mol. Wt	Isotype/Source
ADAM9 (D64B5) Rabbit mAb	4151	20 µl	100-115, 75-80 kDa	Rabbit IgG
DLL1 Antibody	2588	20 µl	82 kDa	Rabbit
DLL3 (G93) Antibody	2483	20 µl	65 kDa	Rabbit
DLL4 Antibody	2589	20 µl	75-80 kDa	Rabbit
Jagged1 (28H8) Rabbit mAb	2620	20 µl	180 kDa	Rabbit IgG
Jagged2 (C23D2) Rabbit mAb	2210	20 µl	150 kDa	Rabbit IgG
Numb (C29G11) Rabbit mAb	2756	20 µl	72, 74 kDa	Rabbit IgG
RBPSUH (D10A4) XP® Rabbit mAb	5313	20 µl	61 kDa	Rabbit IgG
TACE (D22H4) Rabbit mAb	6978	20 µl	135 kDa	Rabbit IgG
Anti-rabbit IgG, HRP-linked Antibody	7074	100 µl		Goat

Please visit cellsignal.com for individual component applications, species cross-reactivity, dilutions, protocols, and additional product information.

Description

The Notch Receptor Interaction Antibody Sampler Kit provides an economical means to evaluate Notch signaling. The kit contains enough primary antibody to perform two western blots per primary.

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Background

Notch signaling is activated upon engagement of the Notch receptor with its ligands, the Delta, Serrate, Lag2 (DSL) single-pass type I membrane proteins. DSL proteins contain multiple EGF-like repeats and a DSL domain that is required for binding to Notch (1,2). Five DSL proteins have been identified in mammals: Jagged1, Jagged2, Delta-like (DLL) 1, 3, and 4 (3). Ligand binding to the Notch receptor results in two sequential proteolytic cleavages of the receptor by the ADAM protease and the γ -secretase complex. The intracellular domain of Notch is released and then translocates to the nucleus where it activates transcription. Notch ligands may also be processed in a similar manner, suggesting bi-directional signaling through receptor-ligand interactions (4-6).

TNF- α converting enzyme (TACE), also known as ADAM17, is a transmembrane metalloprotease that plays a key role in the cleavage of a number cell surface molecules in a process known as "shedding". TACE is abundantly expressed in many adult tissues, but in fetal development, expression is differentially regulated (7). TACE activates Notch in a ligand-independent manner and has been shown to play a role in the development of the *Drosophila* nervous system (8).

Recombining Binding Protein, **S**uppressor of **H**airless (RBPSUH), also termed RBP-J or CSL, is the DNA-binding component of the transcription complex regulated by canonical Notch signaling. In the absence of Notch activation, RBPSUH suppresses target gene expression through interactions with a co-repressor complex containing histone deacetylase. Upon activation of Notch receptors, the Notch intracellular domain (NICD) translocates to the nucleus and binds to RBPSUH. This displaces the co-repressor complex and replaces it with a transcription activation complex that includes Mastermind-like (MAML) proteins and histone acetylase p300, leading to transcriptional activation of Notch target genes (9-11).

Numb contains an amino-terminal phosphotyrosine-binding (PTB) domain and carboxy-terminal endocytic binding motifs for α -adaptin and EH (Eps15 homology) domain-containing proteins, indicating a role in endocytosis (12,13). There are four mammalian Numb splicing isoforms that are differentially expressed and may have distinct functions (14-16). Numb acts as a negative regulator of Notch signaling by promoting ubiquitination and degradation of Notch (17). The protein is asymmetrically segregated into one daughter cell during cell division, producing two daughter cells with different responses to Notch signaling and different cell fates (18,19).

Background References

1. Wilson, A. and Radtke, F. (2006) *FEBS Lett* 580, 2860-8.
2. Hansson, E.M. et al. (2004) *Semin Cancer Biol* 14, 320-8.
3. Chiba, S. (2006) *Stem Cells* 24, 2437-47.
4. Bland, C.E. et al. (2003) *J Biol Chem* 278, 13607-10.
5. Six, E. et al. (2003) *Proc Natl Acad Sci U S A* 100, 7638-43.
6. LaVoie, M.J. and Selkoe, D.J. (2003) *J Biol Chem* 278, 34427-37.
7. Black, R.A. et al. (1997) *Nature* 385, 729-33.
8. Delwig, A. and Rand, M.D. (2008) *Cell Mol Life Sci* 65, 2232-43.
9. Ehebauer, M. et al. (2006) *Sci STKE* 2006, cm7.
10. Borggrefe, T. and Oswald, F. (2009) *Cell Mol Life Sci* 66, 1631-46.
11. Kopan, R. and Ilagan, M.X. (2009) *Cell* 137, 216-33.
12. Berdnik, D. et al. (2002) *Dev Cell* 3, 221-31.
13. Santolini, E. et al. (2000) *J Cell Biol* 151, 1345-52.
14. Dho, S.E. et al. (1999) *J Biol Chem* 274, 33097-104.
15. Verdi, J.M. et al. (1999) *Proc Natl Acad Sci U S A* 96, 10472-6.
16. Verdi, J.M. et al. (1999) *Proc Natl Acad Sci U S A* 96, 10472-6.
17. McGill, M.A. and McGlade, C.J. (2003) *J Biol Chem* 278, 23196-203.
18. Verdi, J.M. et al. (1996) *Curr Biol* 6, 1134-45.
19. Reugels, A.M. et al. (2006) *Dev Dyn* 235, 934-48.

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

U.S. Patent No. 7,429,487, foreign equivalents, and child patents deriving therefrom.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.