20C
e at -
Stor
61
60

AxI (C89E7) Rabbit mAb



877-616-CELL (2355)

orders@cellsignal.com

877-678-TECH (8324) Support:

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

Applications: WB, W-S, IP, IHC-P, IF- IC, FC-FP	Reactivity: H Mk	Sensitivity: Endogenous	MW (kDa): 138	Source/Isotype: Rabbit IgG	UniProt ID: #P30530	Entrez-Gene Id 558	
Product Usage Information	Aį	Application			Dilution		
	W	Western Blotting			1:1000		
	Si	mple Western™		1:10 - 1:50			
	Im	Immunoprecipitation			1:50		
	Im	Immunohistochemistry (Paraffin)			1:300 - 1:1200		
	Im	Immunofluorescence (Immunocytochemistry)			1:100 - 1:400		
	Fl	Flow Cytometry (Fixed/Permeabilized)			1:100 - 1:400		
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μ g/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody.					
	Foi	For a carrier free (BSA and azide free) version of this product see product #80287.					
Specificity / Sensitiv	,	Axl (C89E7) Rabbit mAb detects endogenous levels of total Axl protein and does not cross-react with Tyro3.					
Source / Purification		Monoclonal antibody is produced by immunizing animals with a recombinant human fragment of the Axl protein.					
Background	intr mo is s act (2,7	AxI, Sky, and Mer are three members of a receptor tyrosine kinase (RTK) family that share a conserved intracellular tyrosine kinase domain and an extracellular domain similar to those seen in cell adhesion molecules. These RTKs bind the vitamin K-dependent protein growth-arrest-specific gene 6 (Gas6), which is structurally related to the protein S anticoagulation factor (1). Upon binding to its receptor, Gas6 activates phosphatidylinositol 3-kinase (PI3K) and its downstream targets Akt and S6K, as well as NF-kB (2,3). A large body of evidence supports a role for Gas6/AxI signaling in cell growth and survival in normal and cancer cells (4).					
Background Refere	2. [3. l	 Crosier, K.E. and Crosier, P.S. (1997) Pathology 29, 131-5. Demarchi, F. et al. (2001) J Biol Chem 276, 31738-44. Lee, W.P. et al. (2002) Oncogene 21, 329-36. Bellosta, P. et al. (1997) Oncogene 15, 2387-97. 					
Species Reactivity	Spe	cies reactivity is dete	ermined by testing	g in at least one approve	ed application (e.g., we	estern blot).	

Western Blot Buffer IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS,

0.1% Tween® 20 at 4°C with gentle shaking, overnight.

WB: Western Blotting **W-S:** Simple Western™ **IP:** Immunoprecipitation **Applications Key**

IHC-P: Immunohistochemistry (Paraffin) IF-IC: Immunofluorescence (Immunocytochemistry)

FC-FP: Flow Cytometry (Fixed/Permeabilized)

H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster **Cross-Reactivity Key**

X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse

GP: Guinea Pig Rab: rabbit All: all species expected

Trademarks and **Patents**

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

U.S. Patent No. 7,429,487, foreign equivalents, and child patents deriving therefrom.

Limited Uses

AxI (C89E7) Rabbit mAb (#8661) Datasheet Without Images Cell Signaling Technology All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.