e at -20C	Bid (3C5) Mouse mAb			
Store		Orders:	877-616-CELL (2355) orders@cellsignal.com	
762		Support:	877-678-TECH (8324)	
#876		Web:	info@cellsignal.com cellsignal.com	
#		3 Trask Lane Danvers Ma	assachusetts 01923 USA	

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: WB, IP	Reactivity: H	Sensitivity: Endogenous	MW (kDa): 22	Source/Isotype: Mouse IgG1	UniProt ID: #P55957	Entrez-Gene Id: 637		
Product Usage		Application			Dilution			
Information	١	Western Blotting			1:1000			
	I	Immunoprecipitation			1:50			
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody.						
Specificity / Sensitivity Bid (3C5) Mouse mAb recognizes endogenous levels of total Bid proteir react with endogenous levels of tBid during apoptosis.					d protein. This antibody	/ does not cross-		
Source / Purification		Monoclonal antibody is produced by immunizing animals with a recombinant protein specific to the full- length sequence of human Bid protein.						
Background Background References		Bid is a pro-apoptotic "BH3 domain-only" member of the Bcl-2 family originally discovered to interact with both the anti-apoptotic family member Bcl-2 and the pro-apoptotic protein Bax (1). Bid is normally localized in the cytosolic fraction of cells as an inactive precursor and is cleaved at Asp60 by caspase-8 during Fas signaling, leading to translocation of the carboxyl terminal p15 fragment (tBid) to the mitochondrial outer membrane (2-4). Translocation of Bid is associated with release of cytochrome c from the mitochondria, leading to complex formation with Apaf-1 and caspase-9 and resulting in caspase-9 activation (5-7). Thus, Bid relays an apoptotic signal from the cell surface to the mitochondria triggering caspase activation (8,9).						
		. Wang, K. et al. (1996) . Luo, X. et al. (1998) <i>Ce</i> . Li, H. et al. (1998) <i>Cell</i> . Gross, A. et al. (1999) . Li, P. et al. (1997) <i>Cell</i> . Zou, H. et al. (1999) J . Saleh, A. et al. (1999) . Yin, X.M. et al. (1999) . Korsmeyer, S.J. et al. (ell 94, 481-90. 94, 491-501. J Biol Chem 27 91, 479-89. Biol Chem 274, J Biol Chem 274, Nature 400, 886	4, 1156-63. 11549-56. 4, 17941-5. 5-91.				
Species Reactivity		pecies reactivity is deter	mined by testing	g in at least one approve	ed application (e.g., we	estern blot).		
Western Blot Buffer		IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.						
Applications Key		WB: Western Blotting IP: Immunoprecipitation						
Cross-Reactivity Key		 H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse GP: Guinea Pig Rab: rabbit All: all species expected 						
Trademarks and Patents		Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.						
Limited Uses		Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.						

Bid (3C5) Mouse mAb (#8762) Datasheet Without Images Cell Signaling Technology

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.