

#8779 Store at -20C

## SirT5 (D5E11) Rabbit mAb



**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com

**Support:** 877-678-TECH (8324)

**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

**For Research Use Only. Not for Use in Diagnostic Procedures.**

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
WB	H M Mk	Endogenous	30	Rabbit IgG	#Q9NXA8	23408

<b>Product Usage Information</b>	<b>Application</b> Western Blotting	<b>Dilution</b> 1:1000
<b>Storage</b>	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.	
<b>Specificity / Sensitivity</b>	SirT5 (D5E11) Rabbit mAb recognizes endogenous levels of total SirT5 protein. This antibody does not cross-react with other sirtuin proteins.	
<b>Source / Purification</b>	Monoclonal antibody is produced by immunizing animals with a recombinant protein specific to the full-length human SirT5 protein.	
<b>Background</b>	<p>The Silent Information Regulator (SIR2) family of genes is a highly conserved group of genes that encode nicotinamide adenine dinucleotide (NAD)-dependent protein deacetylases, also known as Class III histone deacetylases. The first discovered and best characterized of these genes is <i>Saccharomyces cerevisiae</i> Sir2, which is involved in silencing of mating type loci, telomere maintenance, DNA damage response, and cell aging (1). SirT5, a mammalian homolog of Sir2, is localized to the mitochondria and has been implicated in the regulation of cell metabolism (2,3). SirT5 deacetylates carbamoyl phosphate synthetase 1 (CPS1) in the mitochondrial matrix and increases its activity in response to fasting, allowing for adaptation to increased amino acid catabolism (4). SirT5 has also been shown to deacetylate cytochrome c in the mitochondrial intermembrane space (5). In addition to its deacetylase activity, SirT5 contains lysine desuccinylase and demalonylase activity (6,7). Succinyl-lysine and malonyl-lysine modifications occur in a variety of organisms and these post-translational modifications are found on many metabolic enzymes (6-8). Like phosphorylation of serine, threonine, and tyrosine residues, lysine succinylation and malonylation induces a change of two negative charges from a +1 to a -1 charge at physiological pH, and are thought to serve similar functions in the regulation of protein activity, protein-protein interactions, and protein stability. SirT5 knockout mice show increased levels of succinyl-lysine and malonyl-lysine protein modifications in the liver, including increased succinylation of CPS1, a known target of SirT5, suggesting that SirT5 functions to regulate metabolic enzymes through its deacetylase, desuccinylase, and demalonylase activities (6,7).</p>	
<b>Background References</b>	<ol style="list-style-type: none"> <li>Guarente, L. (1999) <i>Nat Genet</i> 23, 281-5.</li> <li>Newman, J.C. et al. (2012) <i>J Biol Chem</i> , .</li> <li>He, W. et al. (2012) <i>Trends Endocrinol Metab</i> 23, 467-76.</li> <li>Nakagawa, T. et al. (2009) <i>Cell</i> 137, 560-70.</li> <li>Schlicker, C. et al. (2008) <i>J Mol Biol</i> 382, 790-801.</li> <li>Du, J. et al. (2011) <i>Science</i> 334, 806-9.</li> <li>Peng, C. et al. (2011) <i>Mol Cell Proteomics</i> 10, M111.012658.</li> <li>Zhang, Z. et al. (2011) <i>Nat Chem Biol</i> 7, 58-63.</li> </ol>	

<b>Species Reactivity</b>	Species reactivity is determined by testing in at least one approved application (e.g., western blot).
<b>Western Blot Buffer</b>	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.
<b>Applications Key</b>	<b>WB:</b> Western Blotting
<b>Cross-Reactivity Key</b>	<b>H:</b> human <b>M:</b> mouse <b>R:</b> rat <b>Hm:</b> hamster <b>Mk:</b> monkey <b>Vir:</b> virus <b>Mi:</b> mink <b>C:</b> chicken <b>Dm:</b> D. melanogaster <b>X:</b> Xenopus <b>Z:</b> zebrafish <b>B:</b> bovine <b>Dg:</b> dog <b>Pg:</b> pig <b>Sc:</b> S. cerevisiae <b>Ce:</b> C. elegans <b>Hr:</b> horse <b>GP:</b> Guinea Pig <b>Rab:</b> rabbit <b>All:</b> all species expected

**Trademarks and Patents**

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](https://cellsignal.com/trademarks) for more information.

**Limited Uses**

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.