

**#8886** Store at -20°C

## PICH (D4G8) Rabbit mAb


**Cell Signaling**  
TECHNOLOGY®

**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com

**Support:** 877-678-TECH (8324)

**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

**For Research Use Only. Not for Use in Diagnostic Procedures.**

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
WB	H Mk	Endogenous	175	Rabbit IgG	#Q2NKX8	54821

### Product Usage Information

#### Application

Western Blotting

#### Dilution

1:1000

### Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

### Specificity / Sensitivity

PICH (D4G8) Rabbit mAb recognizes endogenous levels of total PICH protein.

### Species predicted to react based on 100% sequence homology:

Hamster

### Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Gly610 of human PICH protein.

### Background

PICH is a helicase of the SNF2 family of ATPases and is essential for proper chromosome segregation during mitosis (1). While PICH was originally proposed to participate in spindle assembly checkpoint signaling (1), that function was subsequently called into question (2). When phosphorylated at Thr1063 by CDK1, PICH binds the polo-box domain of the mitotic kinase PLK1 (1) and targets it to chromosome arms (3), where it appears to facilitate proper chromosome arm cohesion (4). PICH is also a substrate of PLK1 (1). Localized to the cytoplasm during interphase, PICH begins to accumulate at centromeres and kinetochores in prometaphase (4). As chromosomes begin to separate at the onset of anaphase, PICH associates with ultrafine threads between sister centromeres thought to be composed of entangled DNA (5), a natural consequence of DNA replication. PICH is proposed to cooperate with BLM, a RecQ-like helicase implicated in the genetic disorder Bloom's Syndrome, to displace centromeric histones along these threads, thus enabling them to span large distances without breaking (6). This provides a temporal window for topoisomerase II $\alpha$ -mediated disentanglement (7). Defects in PICH or BLM disrupt proper chromatid segregation and result in the formation of micronuclei (6).

### Background References

1. Baumann, C. et al. (2007) *Cell* 128, 101-14.
2. Hübner, N.C. et al. (2010) *Chromosoma* 119, 149-65.
3. Leng, M. et al. (2008) *Cell Cycle* 7, 1480-9.
4. Kurasawa, Y. and Yu-Lee, L.Y. (2010) *Mol Biol Cell* 21, 1188-99.
5. Wang, L.H. et al. (2008) *Chromosoma* 117, 123-35.
6. Ke, Y. et al. (2011) *EMBO J* 30, 3309-21.
7. Nitiss, J.L. (2009) *Nat Rev Cancer* 9, 327-37.

### Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

### Western Blot Buffer

**IMPORTANT:** For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

### Applications Key

**WB:** Western Blotting

### Cross-Reactivity Key

**H:** human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster  
**X:** Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse  
**GP:** Guinea Pig **Rab:** rabbit **All:** all species expected

## **Trademarks and Patents**

## **Limited Uses**

### **PICH (D4G8) Rabbit mAb (#8886) Datasheet Without Images Cell Signaling Technology**

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](https://cellsignal.com/trademarks) for more information.

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.