EpCAM (VU1D9) Mouse mAb (PE Conjugate)



Orders:

877-616-CELL (2355) orders@cellsignal.com

Support:

877-678-TECH (8324)

Web:

info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only, Not for Use in Diagnostic Procedures.

Applications: FC-FP, FC-L	Reactivity: H	Sensitivity: Endogenous	Source/Isotype: Mouse IgG1	UniProt ID: #P16422	Entrez-Gene Id: 4072	
Product Usage Information	Application			Dilution		
	Flow Cytometry (Fixed/Permeabilized)			1:50 - 1:200		
	Flo	w Cytometry (Live)	1:50 - 1:200		
Storage		Supplied in PBS (pH 7.2), less than 0.1% sodium azide and 2 mg/ml BSA. Store at 4° C. Do not aliquot the antibodies. Protect from light. Do not freeze.				
Specificity / Sensitiv	ity EpC	EpCAM (VU1D9) Mouse mAb (PE Conjugate) detects endogenous levels of total EpC				
Source / Purification	Mor	Monoclonal antibody is produced by immunizing animals with NCI-H69 small cell lung carcinoma cells.				
Product Description	flow	This Cell Signaling Technology antibody is conjugated to phycoerythrin (PE) and tested in-house for direct flow cytometry analysis in human cells. The antibody is expected to exhibit the same species cross-reactivity as the unconjugated EpCAM (VU1D9) Mouse mAb #2929.				
Background	med EpC cell it ha	Epithelial cell adhesion and activating molecule (EpCAM/CD326) is a transmembrane glycoprotein that mediates Ca2+-independent, homophilic adhesions on the basolateral surface of most epithelial cells. EpCAM is not expressed in adult squamous epithelium, but it is highly expressed in adeno and squamous cell carcinomas (1). Research studies identified EpCAM as one of the first tumor-associated antigens, and it has long been a marker of epithelial and tumor tissue. Investigators have shown that EpCAM is highly expressed in cancer cells (reviewed in 2,3).				
Background Referen		•	04) Hum Pathol 35, 122-8. Gires, O. (2007) Br J Cancer 96,	417-23.		

3. Armstrong, A. and Eck, S.L. (2003) Cancer Biol Ther 2, 320-6.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Applications Key

FC-FP: Flow Cytometry (Fixed/Permeabilized) FC-L: Flow Cytometry (Live)

Cross-Reactivity Key

H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse

GP: Guinea Pig Rab: rabbit All: all species expected

Trademarks and **Patents**

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or

EpCAM (VU1D9) Mouse mAb (PE Conjugate) (#8995) Datasheet Without Images Cell Signaling Technology

otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.