3/23/24, 11:36 AM Revision 3

Phospho-p90RSK (Thr359/Ser363) Antibody						Cell Signaling	
Stor					Orders:	877-616-CELL (2355) orders@cellsignal.com	
4					Support:	877-678-TECH (8324)	
#9344					Web:	info@cellsignal.com cellsignal.com	
3 Hask Lane   Danvers   Massachuseus   01923   054							
For Research Use Only.		_					
Applications: WB, IP	Reactivity: H M R Mk	Sensitivity: Endogenous	<b>MW (kDa):</b> 90	Source: Rabbit	<b>UniProt ID:</b> #Q15349, #Q15418	Entrez-Gene Id: 6196, 6195	
Product Usage	Ар	plication			Dilution		
Information	We	estern Blotting		1:1000	1:1000		
	Im	nunoprecipitation			1:50		
Storage	•	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 $\mu$ g/ml BSA and 50% glycerol. Store at – 20°C. Do not aliquot the antibody.					
Specificity / Sensitivity		Phospho-p90RSK (Thr359/Ser363) Antibody detects endogenous levels of p90RSK when phosphorylated at Thr359/Ser363, and singly phosphorylated at Thr359. This antibody does not recognize p90RSK phosphorylated at other sites. It does not cross-react with phospho-RSK2 or -RSK3 at the homologous sites.					
Source / Purification		Polyclonal antibodies are produced by immunizing animals with a synthetic phosphopeptide corresponding to residues surrounding Thr359/Ser363 of rat RSK1. Antibodies are purified by protein A and peptide affinity chromatography.					
Background		The 90 kDa ribosomal S6 kinases (RSK1-4) are a family of widely expressed Ser/Thr kinases characterized by two nonidentical, functional kinase domains (1) and a carboxy-terminal docking site for extracellular signal-regulated kinases (ERKs) (2). Several sites both within and outside of the RSK kinase domain, including Ser380, Thr359, Ser363, and Thr573, are important for kinase activation (3). RSK1-3 are activated via coordinated phosphorylation by MAPKs, autophosphorylation, and phosphoinositide-3-OH kinase (PI3K) in response to many growth factors, polypeptide hormones, and neurotransmitters (3).					
Background References		<ol> <li>Fisher, T.L. and Blenis, J. (1996) Mol Cell Biol 16, 1212-9.</li> <li>Smith, J.A. et al. (1999) J Biol Chem 274, 2893-8.</li> <li>Dalby, K.N. et al. (1998) J Biol Chem 273, 1496-505.</li> </ol>					
Species Reactivity		Species reactivity is determined by testing in at least one approved application (e.g., western blot).					
Western Blot Buffer		IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.					
Applications Key		WB: Western Blotting IP: Immunoprecipitation					
Cross-Reactivity Key		H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse GP: Guinea Pig Rab: rabbit All: all species expected					
Trademarks and Patents		Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.					
Limited Uses		Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.					
		Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the					

Phospho-p90RSK (Thr359/Ser363) Antibody (#9344) Datasheet Without Images Cell Signaling Technology

end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.