Stat6 Antibody



Orders: 877-616-CELL (2355)

orders@cellsignal.com

877-678-TECH (8324) Support:

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

| Applications: WB, IP | Reactivity: H M R | Sensitivity: Endogenous | MW (kDa): 110 | Source: Rabbit | UniProt ID: #P42226 | Entrez-Gene Id 6778 | |
|---|--|--|-------------------------|--------------------------|-------------------------------|------------------------|--|
| Product Usage Information | Ар | plication | | | Dilution | | |
| | We | Western Blotting | | | 1:1000 | | |
| | Imr | munoprecipitation | | 1:100 | | | |
| Storage | Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μ g/ml BSA and 50% glycerol. Store at – 20°C. Do not aliquot the antibody. | | | | | glycerol. Store at – | |
| Specificity / Sensitivity | | Stat6 Antibody detects endogenous levels of total Stat6 protein. | | | | | |
| Species predicted to react based on 100% sequence homology: | | Bovine | | | | | |
| Source / Purification | | Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues surrounding amino acid 620 of human Stat6. Antibodies are purified by protein A and peptide affinity chromatography. | | | | | |
| Background | | Upon activation by Janus kinases, Stat6 translocates to the nucleus where it regulates cytokine-induced gene expression. Stat6 is activated via phosphorylation at Tyr641 and is required for responsiveness to IL-4 and IL-13 (1-4). In addition, Stat6 is activated by IFN- α in B cells, where it forms transcriptionally active complexes with Stat2 and p48 (5,6). Protein phosphatase 2A is also involved in regulation of IL-4-mediated Stat6 signaling (7). | | | | | |

Background References

- 1. Nelms, K. et al. (1999) Ann. Rev. Immunol. 17, 701-738.
- 2. Malabarba, M.G. et al. (1996) Biochem. J. 319, 865-872.
- 3. Hou, J. et al. (1994) Science 265, 1701-1706.
- 4. Quelle, F.W. et al. (1995) Mol. Cell. Biol. 15, 3336-3343.
- 5. Takeda, K. et al. (1996) Nature 380, 627-630.
- 6. Gupta, S. et al. (1999) J. Immunol. 163, 3834-3841.
- 7. Woetmann, A. et al. (2003) J. Biol. Chem. 278, 2787-2791.

Species reactivity is determined by testing in at least one approved application (e.g., western blot). **Species Reactivity**

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, Western Blot Buffer

0.1% Tween® 20 at 4°C with gentle shaking, overnight.

WB: Western Blotting IP: Immunoprecipitation **Applications Key**

H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster **Cross-Reactivity Key**

X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse

GP: Guinea Pig Rab: rabbit All: all species expected

Trademarks and **Patents**

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more

information.

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the **Limited Uses**

following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in

Stat6 Antibody (#9362) Datasheet Without Images Cell Signaling Technology writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.