Conjugate)

Store at +4C

9488

Phospho-p44/42 MAPK (Erk1/2) (Thr202/Tyr204) (D13.14.4E) XP[®]



Orders:	orders@cellsignal.com
Support:	877-678-TECH (8324)
Web:	info@cellsignal.com cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use	Only. Not for	Use in Diagnosti	c Procedures.

Rabbit mÁb (Magnetic Bead

Applications: IP	Reactivity: H M R Hm Mk Mi Dm Z B Dg Pg Sc	Sensitivity: Endogenous	MW (kDa): 44, 42	Source/Isotype: Rabbit IgG	UniProt ID: #P27361, #P28482	Entrez-Gene Id: 5595, 5594	
Product Usage	Ар	plication			Dilution		
Information	Im	munoprecipitation			1:20		
Storage	Sup	plied in PBS Buffer	(pH 7.2), 0.1% T	Tween $^{\circledast}$ 20. Store at 4°C. Do not aliquot the antibodies.			
Specificity / Sensitivity		Phospho-p44/42 MAPK (Erk1/2) (Thr202/Tyr204) (D13.14.4E) XP [®] Rabbit mAb detects endogenous levels of p44 and p42 MAP Kinase (Erk1 and Erk2) when dually phosphorylated at Thr202 and Tyr204 of Erk1 (Thr185 and Tyr187 of Erk2), and singly phosphorylated at Thr202 of Erk1 (Thr185 of Erk2). This antibody does not cross-react with the corresponding phosphorylated residues of either JNK/SAPK or p38 MAP kinases.					
Species predicte react based on 1 sequence homol	.00%	cken, C. elegans					
Source / Purifica		Monoclonal antibody is produced by immunizing animals with a synthetic phosphopeptide corresponding to residues surrounding Thr202/Tyr204 of human p44 MAP kinase.					
Product Descrip	moo (Th	This Cell Signaling Technology antibody is immobilized by the covalent reaction of hydrazinonicotinamide- modifed antibody with formylbenzamide-modified magnetic bead. Phospho-p44/42 MAPK (Erk1/2) (Thr202/Tyr204) (D13.14.4E) XP [®] Rabbit mAb (Magnetic Bead Conjugate) is useful for immunoprecipitation of phosphorylated Erk protein.					
MW (kDa)				44, 42			
Background	kina The extr con thre MA hav are pho dow (8,9	Mitogen-activated protein kinases (MAPKs) are a widely conserved family of serine/threonine protein kinases involved in many cellular programs, such as cell proliferation, differentiation, motility, and death. The p44/42 MAPK (Erk1/2) signaling pathway can be activated in response to a diverse range of extracellular stimuli, including mitogens, growth factors, and cytokines (1-3), and research investigators consider it an important target in the diagnosis and treatment of cancer (4). Upon stimulation, a sequential three-part protein kinase cascade is initiated, consisting of a MAP kinase kinase (MAPKK or MAP3K), a MAP kinase kinase (MAPKK or MAP3K), a MAP kinase kinase (MAPKK or MAP2K), and a MAP kinase (MAPK). Multiple p44/42 MAP3Ks have been identified, including members of the Raf family, as well as Mos and Tpl2/COT. MEK1 and MEK2 are the primary MAPKKs in this pathway (5,6). MEK1 and MEK2 activate p44 and p42 through phosphorylation of activation loop residues Thr202/Tyr204 and Thr185/Tyr187, respectively. Several downstream targets of p44/42 have been identified, including p90RSK (7) and the transcription factor Elk-1 (8,9). p44/42 are negatively regulated by a family of dual-specificity (Thr/Tyr) MAPK phosphatases, known as DUSPs or MKPs (10), along with MEK inhibitors, such as U0126 and PD98059.			tility, and death. range of ch investigators ation, a sequential (MAPKKK or e p44/42 MAP3Ks . MEK1 and MEK2 mough rely. Several cription factor Elk-1		

1/1/24, 11 Back	:10 AM Phospho ground References	 -p44/42 MAPK (Erk1/2) (Thr202/Tyr204) (D13.14.4E) XP® Rabbit mAb (Magnetic Bead Conjugate) 1. Roux, P.P. and Blenis, J. (2004) <i>Microbiol Mol Biol Rev</i> 68, 320-44. 2. Baccarini, M. (2005) <i>FEBS Lett</i> 579, 3271-7. 3. Meloche, S. and Pouysségur, J. (2007) <i>Oncogene</i> 26, 3227-39. 4. Roberts, P.J. and Der, C.J. (2007) <i>Oncogene</i> 26, 3291-310. 5. Rubinfeld, H. and Seger, R. (2005) <i>Mol Biotechnol</i> 31, 151-74. 6. Murphy, L.O. and Blenis, J. (2006) <i>Trends Biochem Sci</i> 31, 268-75. 7. Dalby, K.N. et al. (1998) <i>J Biol Chem</i> 273, 1496-505. 8. Marais, R. et al. (1993) <i>Cell</i> 73, 381-93. 9. Kortenjann, M. et al. (1994) <i>Mol Cell Biol</i> 14, 4815-24. 10. Owens, D.M. and Keyse, S.M. (2007) <i>Oncogene</i> 26, 3203-13.
Spec	ies Reactivity	Species reactivity is determined by testing in at least one approved application (e.g., western blot).
Appli	ications Key	IP: Immunoprecipitation
Cros	s-Reactivity Key	H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse GP: Guinea Pig Rab: rabbit All: all species expected
Trade Pater	emarks and nts	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. XP is a registered trademark of Cell Signaling Technology, Inc. All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.
Limit	ed Uses	Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.
		Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products for the purpose of developing any products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.