Store at -20C

Cleaved PARP (Asp214) Antibody



Orders:

877-616-CELL (2355) orders@cellsignal.com

Support:

877-678-TECH (8324)

Web:

info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

Applications: WB	Reactivity: R	Sensitivity: Endogenous	MW (kDa): 89	Source: Rabbit	UniProt ID: #P09874	Entrez-Gene Id 142	
Product Usage Information	Application			Dilution			
	We	stern Blotting			1:1000		
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μ g/ml BSA and 50% glycerol. Store at $-$ 20°C. Do not aliquot the antibody.					
Specificity / Sensitiv	resu	Cleaved PARP (Asp214) Antibody detects endogenous levels of the large fragment of PARP1 (89 kDa) resulting from caspase cleavage. The antibody does not recognize full length PARP1 or other PARP isoforms.					
Source / Purification	carb	Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to carboxy-terminal residues surrounding Asp214 in rat PARP. Antibodies are purified by protein A and peptide affinity chromatography.					
Background	to el one Asp. carb	PARP, a 116 kDa nuclear poly (ADP-ribose) polymerase, appears to be involved in DNA repair in response to environmental stress (1). This protein can be cleaved by many ICE-like caspases <i>in vitro</i> (2,3) and is one of the main cleavage targets of caspase-3 <i>in vivo</i> (4,5). In human PARP, the cleavage occurs between Asp214 and Gly215, which separates the PARP amino-terminal DNA-binding domain (24 kDa) from the carboxy-terminal catalytic domain (89 kDa) (2,4). PARP helps cells to maintain their viability; cleavage of PARP facilitates cellular disassembly and serves as a marker of cells undergoing apoptosis (6).					
	(Thi	(This product is sold under license from Promega Corp., U.S. Patent No. 6,350,452.)					
Background Refere	2. La	 Satoh, M.S. and Lindahl, T. (1992) <i>Nature</i> 356, 356-358. Lazebnik, Y. A. et al. (1994) <i>Nature</i> 371, 346-347. Cohen, G.M. (1997) <i>Biochem. J.</i> 326, 1-16. 					

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry

milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting

Cross-Reactivity Key

H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster

X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse

GP: Guinea Pig Rab: rabbit All: all species expected

4. Nicholson, D. W. et al. (1995) Nature 376, 37-43. 5. Tewari, M. et al. (1995) Cell 81, 801-809.

6. Oliver, F.J. et al. (1998) J. Biol. Chem. 273, 33533-33539.

Trademarks and **Patents**

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Cleaved PARP (Asp214) Antibody (#9545) Datasheet Without Images Cell Signaling Technology

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.