Store at -200

Phospho-PTEN (Ser380) Antibody



Orders: 877-616-CELL (2355)

orders@cellsignal.com

877-678-TECH (8324) Support:

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

Applications: WB, IP	Reactivity: H M R	Sensitivity: Endogenous	MW (kDa): 54	Source: Rabbit	UniProt ID: #P60484	Entrez-Gene Id 5728	
Product Usage Information	Ар	Application			Dilution		
	We	Western Blotting			1:1000		
	lmı	Immunoprecipitation			1:50		
Storage		plied in 10 mM sodi C. Do not aliquot the		i), 150 mM NaCl, 10	00 μg/ml BSA and 50% ç	ylycerol. Store at –	
Specificity / Sensitivity Phospho-PTEN (Ser380) Antibody d Ser380.		O) Antibody detects	cts endogenous levels of PTEN only when phosphorylated at				
Species predicted react based on 10 sequence homological	0%	cken					

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic phosphopeptide corresponding to residues surrounding Ser380 of human PTEN. Antibodies are purified by protein A and peptide affinity chromatography.

Background

PTEN (phosphatase and tensin homologue deleted on chromosome ten), also referred to as MMAC (mutated in multiple advanced cancers) phosphatase, is a tumor suppressor implicated in a wide variety of human cancers (1). PTEN encodes a 403 amino acid polypeptide originally described as a dual-specificity protein phosphatase (2). The main substrates of PTEN are inositol phospholipids generated by the activation of the phosphoinositide 3-kinase (PI3K) (3). PTEN is a major negative regulator of the PI3K/Akt signaling pathway (1,4,5). PTEN possesses a carboxy-terminal, noncatalytic regulatory domain with three phosphorylation sites (Ser380, Thr382, and Thr383) that regulate PTEN stability and may affect its biological activity (6,7). PTEN regulates p53 protein levels and activity (8) and is involved in G proteincoupled signaling during chemotaxis (9,10).

Background References

- 1. Cantley, L.C. and Neel, B.G. (1999) Proc Natl Acad Sci USA 96, 4240-5.
- 2. Myers, M.P. et al. (1997) Proc Natl Acad Sci USA 94, 9052-7.
- 3. Myers, M.P. et al. (1998) Proc Natl Acad Sci USA 95, 13513-8.
- 4. Wan, X. and Helman, L.J. (2003) Oncogene 22, 8205-11.
- 5. Wu, X. et al. (1998) Proc Natl Acad Sci USA 95, 15587-91.
- 6. Vazquez, F. et al. (2000) Mol Cell Biol 20, 5010-8.
- 7. Torres, J. and Pulido, R. (2001) J Biol Chem 276, 993-8.
- 8. Freeman, D.J. et al. (2003) Cancer Cell 3, 117-30.
- 9. Funamoto, S. et al. (2002) Cell 109, 611-23.
- 10. lijima, M. and Devreotes, P. (2002) Cell 109, 599-610.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting IP: Immunoprecipitation

Cross-Reactivity Key

Phospho-PTEN (Ser380) Antibody (#9551) Datasheet Without Images Cell Signaling Technology

H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse GP: Guinea Pig Rab: rabbit All: all species expected

Trademarks and Patents

Limited Uses

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.
All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.