#9631 Store at -20C

Phospho-(Ser/Thr) Phe Antibody



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:Reactivity:Sensitivity:Source:WB, IP, E-PAllEndogenousRabbit

Product Usage
InformationApplicationDilutionWestern Blotting1:1000Immunoprecipitation1:100Peptide ELISA (DELFIA)1:500

Storage Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at –

20°C. Do not aliquot the antibody.

Specificity / Sensitivity Phospho-(Ser/Thr) Phe Antibody detects phospho-serine or threonine in the context of tyrosine, tryptophan

or phenylalanine at the -1 position or phenylalanine at the +1 position. The antibody does not cross-react

with the nonphosphorylated form of these motifs, nor does it cross-react with other phosphoserine/threonine-containing proteins and peptides. (U.S. Patent No's.: 6,441,140; 6,982,318; 7,259,022;

7,344,714; U.S.S.N. 11,484,485; and all foreign equivalents.)

Source / Purification Polyclonal antibodies are produced by immunizing animals with a synthetic phospho-serine/threonine-

phenylalanine-containing peptide. Antibodies are purified by protein A and peptide affinity chromatography.

Background A hallmark of signal transduction pathways is the reversible phosphorylation of serine and threonine

residues within specific sequences, or motifs, in target proteins. Specific signaling motifs include not only sequences that are recognized by protein kinases (1), but also those that are recognized by

phosphorylation-dependent binding proteins such as 14-3-3 (2). These modular phosphoprotein interacting domains are critical elements in modulating, directing and amplifying intracellular communications. CST has pioneered the development of phospho-motif specific antibodies, which are invaluable tools for probing

the complexity of phospho-regulatory pathways.

Many critical protein kinases can be regulated by phosphorylation at a specific serine or threonine surrounded by phenylalanine or tyrosine. For example, Akt, a kinase that regulates cell survival, is activated by phosphorylation at Ser473, a site surrounded by phenylalanine and tyrosine (3). RSK1, p70S6K, and certain PKC isoforms also contain a similar consensus phosphorylation site. Phosphorylation of these sites is required for kinase activity (4,5). The Phospho-(Ser/Thr) Phe Antibody is a powerful tool

for discovery of new proteins containing this important regulatory motif.

Background References 1. Pinna, L.A. and Ruzzene, M. (1996) *Biochim Biophys Acta* 1314, 191-225.

2. Yaffe, M.B. and Elia, A.E. (2001) Curr Opin Cell Biol 13, 131-8.

3. Alessi, D.R. et al. (1996) *EMBO J* 15, 6541-51.

4. Dalby, K.N. et al. (1998) J Biol Chem 273, 1496-505.

5. Keranen, L.M. et al. (1995) Curr Biol 5, 1394-1403.

Species Reactivity Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS,

0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key WB: Western Blotting IP: Immunoprecipitation E-P: Peptide ELISA (DELFIA)

Cross-Reactivity Key H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster

X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse

GP: Guinea Pig Rab: rabbit All: all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

Phospho-(Ser/Thr) Phe Antibody (#9631) Datasheet Without Images Cell Signaling Technology

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.