#9681 Store at -20C

Acetylated-Lysine (Ac-K-103) Mouse mAb



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:Reactivity:Sensitivity:Source/Isotype:WB, E-PAllEndogenousMouse IgG2a

 Product Usage Information
 Application
 Dilution

 Western Blotting
 1:1000

 Peptide ELISA (DELFIA)
 1:1000

Storage Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than

0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Specificity / Sensitivity

Acetylated-Lysine (Ac-K-103) Mouse mAb detects proteins only when posttranslationally modified by acetylation on the epsilon-amine groups of lysine residues. Detection of acetylated lysine by this antibody is largely independent of surrounding amino acid sequence. The antibody has been shown to recognize acetylated proteins including histones, p53, CBP, PCAF and chemically acetylated BSA. (U.S. Patent No's.: 6,441,140; 6,982,318; 7,259,022; 7,344,714; U.S.S.N. 11,484,485; and all foreign equivalents.)

Source / PurificationMonoclonal antibody is produced by immunizing animals with a synthetic acetylated lysine-containing peptide.

BackgroundAcetylation of lysine, like phosphorylation of serine, threonine or tyrosine, is an important reversible modification controlling protein activity. The conserved amino-terminal domains of the four core histones

(H2A, H2B, H3, and H4) contain lysines that are acetylated by histone acetyltransferases (HATs) and deacetylated by histone deacetylated by histone deacetylation of histones, transcription factors, and other proteins affects a diverse array of cellular processes including chromatin structure and gene activity, cell growth, differentiation, and apoptosis (2-6). Recent proteomic surveys suggest that acetylation of lysine residues may be a widespread and important form of post-translational protein modification that affects thousands of proteins involved in control of cell cycle and metabolism, longevity, actin polymerization, and nuclear transport (7,8). The regulation of protein acetylation status is impaired in cancer and polyglutamine diseases (9), and HDACs have become

promising targets for anti-cancer drugs currently in development (10).

Background References 1. Hassig, C.A. and Schreiber, S.L. (1997) *Curr Opin Chem Biol* 1, 300-8.

2. Allfrey, V.G. et al. (1964) Proc Natl Acad Sci USA 51, 786-94.

3. Liu, L. et al. (1999) Mol Cell Biol 19, 1202-9.

4. Boyes, J. et al. (1998) Nature 396, 594-8.

5. Polevoda, B. and Sherman, F. (2002) Genome Biol 3, reviews 0006.

6. Yoshida, M. et al. (2003) Prog Cell Cycle Res 5, 269-78.

7. Kim, S.C. et al. (2006) Mol Cell 23, 607-18.

8. Choudhary, C. et al. (2009) Science 325, 834-40.

9. Hughes, R.E. (2002) Curr Biol 12, R141-3.

10. Vigushin, D.M. and Coombes, R.C. (2004) Curr Cancer Drug Targets 4, 205-18.

Species Reactivity Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS,

0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key WB: Western Blotting E-P: Peptide ELISA (DELFIA)

Cross-Reactivity Key H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster

X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse

GP: Guinea Pig Rab: rabbit All: all species expected

4/23/24, 10:30 AM

Trademarks and Patents

Limited Uses

Acetylated-Lysine (Ac-K-103) Mouse mAb (#9681) Datasheet Without Images Cell Signaling Technology

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.
All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.