

#9869 Store at -20°C

Cyclin Antibody Sampler Kit

1 Kit (8 x 20 microliters)



Cell Signaling
TECHNOLOGY®

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Product Includes	Product #	Quantity	Mol. Wt	Isotype/Source
Cyclin A2 (BF683) Mouse mAb	4656	20 µl	55 kDa	Mouse IgE
Cyclin B1 Antibody	4138	20 µl	55 kDa	Rabbit
Cyclin D1 (92G2) Rabbit mAb	2978	20 µl	36 kDa	Rabbit IgG
Cyclin D2 (D52F9) Rabbit mAb	3741	20 µl	31 kDa	Rabbit IgG
Cyclin D3 (DCS22) Mouse mAb	2936	20 µl	31 kDa	Mouse IgG1
Cyclin E1 (HE12) Mouse mAb	4129	20 µl	48 kDa	Mouse IgG1
Cyclin E2 Antibody	4132	20 µl	48 kDa	Rabbit
Cyclin H Antibody	2927	20 µl	36 kDa	Rabbit
Anti-rabbit IgG, HRP-linked Antibody	7074	100 µl		Goat
Anti-mouse IgG, HRP-linked Antibody	7076	100 µl		Horse

Please visit cellsignal.com for individual component applications, species cross-reactivity, dilutions, protocols, and additional product information.

Description

The Cyclin Antibody Sampler Kit provides an economical means of evaluating the presence of cyclin proteins in cells. The kit contains enough primary and secondary antibodies to perform two western blot experiments with each antibody.

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Background

Control of the cell cycle is regulated by a multitude of cellular events and processes. The cyclin dependent kinases (CDK) regulate many of these pathways and constitute an active complex when associated with their cyclin partners. This activity is controlled primarily by phosphorylation, which determines subcellular localization of the CDK/cyclin complex (1,2). Some phosphorylation events control the function of cytoplasmic retention sequences while other events regulate nuclear localization and export sequence function (3,4). Cyclin and cyclin-dependent kinase inhibitor (CKI) levels are regulated by ubiquitination and degradation via the ubiquitin proteasome pathway (5). A variety of CKI proteins associate with these complexes and modulate access to regulatory domains on cyclins (6). Additional complexity is generated as the controlled protein levels of each cyclin oscillate with the stages of cell cycle. Increased expression of Cyclin D1 is associated with certain types of cancer (7,8) and may associate with TSC2 (tuberin) independent of its Cdk partner (9).

Background References

1. Kremler, A. et al. (2005) *Cell Mol Life Sci* 62, 1379-87.
2. Toyoshima, F. et al. (1998) *EMBO J* 17, 2728-35.
3. Porter, L.A. and Donoghue, D.J. (2003) *Prog Cell Cycle Res* 5, 335-47.
4. Benzeno, S. et al. (2006) *Oncogene* 25, 6291-303.
5. Diehl, J.A. et al. (1997) *Genes Dev* 11, 957-72.
6. Hirai, H. et al. (1995) *Mol Cell Biol* 15, 2672-81.
7. Sherr, C.J. (1996) *Science* 274, 1672-7.
8. Kim, J.K. and Diehl, J.A. (2009) *J Cell Physiol* 220, 292-6.
9. Zacharek, S.J. et al. (2005) *Cancer Res* 65, 11354-60.

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.
All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in

writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.