

Store at -20C
#9893

PZR (D17B10) XP® Rabbit mAb



Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
WB, IP, IF-IC	H	Endogenous	30-50	Rabbit IgG	#O95297	9019

Product Usage Information

Application

Western Blotting
Immunoprecipitation
Immunofluorescence (Immunocytochemistry)

Dilution

1:1000
1:50
1:50

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Specificity / Sensitivity

PZR (D17B10) XP® Rabbit mAb recognizes endogenous levels of total PZR protein.

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Leu209 of human PZR protein.

Background

PZR (Protein zero related) is an immunoglobulin superfamily protein that specifically binds the tyrosine phosphatase SHP-2 through its intracellular immunoreceptor tyrosine-based inhibitory motifs (ITIMs) (1,2). PZR is phosphorylated by c-Src, c-Fyn, c-Lyn, Csk, and c-Abl (3). PP1, a Src family kinase inhibitor, inhibits PZR phosphorylation (4,5). There are three alternatively spliced isoforms, designated as PZR, PZRα, and PZRβ; both PZRα and PZRβ lack ITIMs (6,7). PZR is the main receptor of ConA and has an important role in cell signaling via c-Src (4). PZR is expressed in many cell types and is localized to cell contacts and intracellular granules in BAECs and mesothelioma (REN) cells. PZR has been implicated as a cell adhesion protein that may be involved in SHP-2-dependent signaling at interendothelial cell contacts (3). Hypertyrosine phosphorylation of PZR was observed during embryogenesis in a mouse model of Noonan syndrome (8).

Background References

1. Zhao, Z.J. and Zhao, R. (1998) *J Biol Chem* 273, 29367-72.
2. Zhao, R. and Zhao, Z.J. (2000) *J Biol Chem* 275, 5453-9.
3. Kusano, K. et al. *Endothelium* 15, 127-36.
4. Zhao, R. et al. (2002) *J Biol Chem* 277, 7882-8.
5. Zhao, R. et al. (2003) *J Biol Chem* 278, 42893-8.
6. Zannettino, A.C. et al. (2003) *Biochem J* 370, 537-49.
7. Zhao, R. and Zhao, Z.J. (2003) *Biochem Biophys Res Commun* 303, 1028-33.
8. Eminaga, S. and Bennett, A.M. (2008) *J Biol Chem* 283, 15328-38.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting **IP:** Immunoprecipitation **IF-IC:** Immunofluorescence (Immunocytochemistry)

Cross-Reactivity Key

H: human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster
X: Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse
GP: Guinea Pig **Rab:** rabbit **All:** all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.
XP is a registered trademark of Cell Signaling Technology, Inc.
All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.