

#9925 Store at -20°C

Phospho-Syk Antibody Sampler Kit

1 Kit (4 x 20 microliters)



Cell Signaling
TECHNOLOGY®

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Product Includes	Product #	Quantity	Mol. Wt	Isotype/Source
Phospho-Syk (Tyr323) Antibody	2715	20 µl	72 kDa	Rabbit
Phospho-Syk (Tyr525/526) (C87C1) Rabbit mAb	2710	20 µl	72 kDa	Rabbit IgG
Phospho-Zap-70 (Tyr319)/Syk (Tyr352) Antibody	2701	20 µl	70 Zap-70, 72 Syk kDa	Rabbit
Syk (D3Z1E) XP® Rabbit mAb	13198	20 µl	72 kDa	Rabbit IgG
Anti-rabbit IgG, HRP-linked Antibody	7074	100 µl		Goat

Please visit cellsignal.com for individual component applications, species cross-reactivity, dilutions, protocols, and additional product information.

Description

Phospho-Syk Sampler Kit provides an economical means to evaluate the activation status of Syk, including the phosphorylation of Tyr323, Tyr352 and Tyr525/526. The control Syk Antibody is also included. The kit contains enough primary and secondary antibodies for two Western blot experiments.

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Background

Syk is a protein tyrosine kinase that plays an important role in intracellular signal transduction in hematopoietic cells (1-3). Syk interacts with immunoreceptor tyrosine-based activation motifs (ITAMs) located in the cytoplasmic domains of immune receptors (4). It couples the activated immunoreceptors to downstream signaling events that mediate diverse cellular responses, including proliferation, differentiation, and phagocytosis (4). There is also evidence of a role for Syk in nonimmune cells and investigators have indicated that Syk is a potential tumor suppressor in human breast carcinomas (5). Tyr323 is a negative regulatory phosphorylation site within the SH2-kinase linker region in Syk. Phosphorylation at Tyr323 provides a direct binding site for the TKB domain of Cbl (6,7). Tyr352 of Syk is involved in the association of PLCγ1 (8). Tyr525 and Tyr526 are located in the activation loop of the Syk kinase domain; phosphorylation at Tyr525/526 of human Syk (equivalent to Tyr519/520 of mouse Syk) is essential for Syk function (9).

Background References

1. Cheng, A.M. and Chan, A.C. (1997) *Curr Opin Immunol* 9, 528-33.
2. Kurosaki, T. (1997) *Curr Opin Immunol* 9, 309-18.
3. Chu, D.H. et al. (1998) *Immunol Rev* 165, 167-80.
4. Turner, M. et al. (2000) *Immunol Today* 21, 148-54.
5. Coopman, P.J. et al. (2000) *Nature* 406, 742-7.
6. Deckert, M. et al. (1998) *J Biol Chem* 273, 8867-74.
7. Rao, N. et al. (2001) *EMBO J* 20, 7085-95.
8. Law, C.L. et al. (1996) *Mol Cell Biol* 16, 1305-15.
9. Zhang, J. et al. (2000) *J Biol Chem* 275, 35442-7.

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. U.S. Patent No. 7,429,487, foreign equivalents, and child patents deriving therefrom. All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the

end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.