

#9935 Store at -20°C

Src Antibody Sampler Kit

1 Kit (5 x 20 microliters)



Cell Signaling
TECHNOLOGY®

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Product Includes	Product #	Quantity	Mol. Wt	Isotype/Source
Phospho-Src Family (Tyr416) (D49G4) Rabbit mAb	6943	20 µl	60 kDa	Rabbit IgG
Non-phospho-Src (Tyr527) Antibody	2107	20 µl	60 kDa	Rabbit
Non-phospho-Src (Tyr416) (7G9) Mouse mAb	2102	20 µl	60 kDa	Mouse IgG2b
Phospho-Src (Tyr527) Antibody	2105	20 µl	60 kDa	Rabbit
Src (32G6) Rabbit mAb	2123	20 µl	60 kDa	Rabbit IgG
Anti-rabbit IgG, HRP-linked Antibody	7074	100 µl		Goat
Anti-mouse IgG, HRP-linked Antibody	7076	100 µl		Horse

Please visit [cellsignal.com](https://www.cellsignal.com) for individual component applications, species cross-reactivity, dilutions, protocols, and additional product information.

Description

The Src Antibody Sampler kit provides an economical means of evaluating total Src protein levels and its phosphorylation status. The kit contains enough primary and secondary antibodies to perform two Western blot experiments per primary antibody.

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Background

The Src family of protein tyrosine kinases, which includes Src, Lyn, Fyn, Yes, Lck, Blk, and Hck, are important in the regulation of growth and differentiation of eukaryotic cells (1). Src activity is regulated by tyrosine phosphorylation at two sites, but with opposing effects. While phosphorylation at Tyr416 in the activation loop of the kinase domain upregulates enzyme activity, phosphorylation at Tyr527 in the carboxy-terminal tail by Csk renders the enzyme less active (2).

Lyn is a member of the Src family that is predominantly expressed in hematopoietic cells (3). Lyn participates in signaling from multiple cell surface receptors such as the B cell Ag receptor (BCR) and CD40 (4).

Lck is essential for T-lymphocyte activation and differentiation (5,6). The activity of Lck is regulated by protein kinases and phosphatases. Phosphorylation of the C-terminal tyrosine 505 serves to downregulate Lck catalytic activity, while phosphorylation at tyrosine 394 leads to an increase in Lck activity (7).

Background References

1. Thomas, S.M. and Brugge, J.S. (1997) *Annu Rev Cell Dev Biol* 13, 513-609.
2. Hunter, T. (1987) *Cell* 49, 1-4.
3. Yamanashi, Y. et al. (1989) *Proc Natl Acad Sci U S A* 86, 6538-42.
4. Yamanashi, Y. et al. (1991) *Science* 251, 192-4.
5. Molina, T.J. et al. (1992) *Nature* 357, 161-4.
6. Straus, D.B. and Weiss, A. (1992) *Cell* 70, 585-93.
7. Chow, L.M. et al. (1993) *Nature* 365, 156-60.

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](https://www.cellsignal.com/trademarks) for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or

therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.