1959 Store at -200

MAML1 (D3E9) Rabbit mAb



Orders: 877-616-CELL (2355)

orders@cellsignal.com

877-678-TECH (8324) Support:

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.							
Applications: WB, IP, ChIP, ChIP-seq	Reactivity: H M R Mk	Sensitivity: Endogenous	MW (kDa): 130	Source/Isotype: Rabbit IgG	UniProt ID: #Q92585	Entrez-Gene Id: 9794	
Product Usage Information				use 5 µl of antibody and dated using SimpleChIF			
	Ap	Application			Dilution		
	We	Western Blotting			1:1000		
	Imr	Immunoprecipitation			1:50		
	Chr	Chromatin IP			1:100		
	Chr	Chromatin IP-seq			1:100		
Storage	• •	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μ g/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20° C. Do not aliquot the antibody.					
Specificity / Sensiti		MAML1 (D3E9) Rabbit mAb recognizes endogenous levels of total MAML1 protein. It does not cross-react with MAML2 or MAML3 protein.					
Source / Purificatio		Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues near the carboxy terminus of human MAML1 protein.					
Background	com com regu (ME carc	Mastermind-like (MAML) family of proteins are homologs of <i>Drosophila</i> Mastermind. The family is composed of three members in mammals: MAML1, MAML2, and MAML3 (1,2). MAML proteins form complexes with the intracellular domain of Notch (ICN) and the transcription factor CSL (RBP-Jk) to regulate Notch target gene expression (3-5). MAML1 also interacts with myocyte enhancer factor 2C (MEF2C) to regulate myogenesis (6). MAML2 is frequently found to be fused with Mucoepidermoid carcinoma translocated gene 1 (MECT1, also know as WAMTP1 or TORC1) in patients with mucoepidermoid carcinomas and Warthin's tumors (7).					
Background Refere	2. Li	 Wu, L. et al. (2002) Mol. Cell Biol. 22, 7688-7700. Lin, S.E. et al. (2002) J. Biol. Chem. 277, 50612-50620. Kitagawa, M. et al. (2001) Mol. Cell Biol. 21, 4337-4346. 					

- 4. Nam, Y. et al. (2003) J. Biol. Chem. 278, 21232-21239.
- 5. Nam, Y. et al. (2006) Cell 124, 973-983.
- 6. Shen, H. et al. (2006) Genes Dev. 20, 675-688.
- 7. Tonon, G. et al. (2003) Nat. Genet. 33, 208-213.

Species Reactivity Species reactivity is determined by testing in at least one approved application (e.g., western blot).

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, Western Blot Buffer

0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting IP: Immunoprecipitation ChIP: Chromatin IP ChIP-seq: Chromatin IP-seq

Cross-Reactivity Key H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster

X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse

GP: Guinea Pig Rab: rabbit All: all species expected

Trademarks and **Patents**

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. SimpleChIP is a registered trademark of Cell Signaling Technology, Inc.

XP is a registered trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more

1/1/24, 12:12 PM **Limited Uses**

MAML1 (D3E9) Rabbit mAb (#11959) Datasheet Without Images Cell Signaling Technology

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.