

**#12402** Store at -20°C

## PI3 Kinase Class II $\alpha$ (D3Q5B) Rabbit mAb


**Cell Signaling**  
TECHNOLOGY®

**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com

**Support:** 877-678-TECH (8324)

**Web:** info@cellsignal.com  
cellsignal.com

**3 Trask Lane | Danvers | Massachusetts | 01923 | USA**
**For Research Use Only. Not for Use in Diagnostic Procedures.**

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
WB, IP	H M R Mk	Endogenous	180	Rabbit IgG	#O00443	5286

### Product Usage Information

#### Application

Western Blotting  
Immunoprecipitation

#### Dilution

1:1000  
1:50

### Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100  $\mu$ g/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at  $-20^{\circ}\text{C}$ . Do not aliquot the antibody.

### Specificity / Sensitivity

PI3 Kinase Class II  $\alpha$  (D3Q5B) Rabbit mAb recognizes endogenous levels of total PI3K class II  $\alpha$  protein.

### Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Gly717 of human PI3K class II  $\alpha$  protein.

### Background

Class II phosphatidylinositol 3-kinases (PI3K) contain a C-terminal C2 domain that is unique to the class II isoforms of the PI3K family. This C2 domain mediates protein and phospholipid binding activities (1,2). PI3K Class II  $\alpha$  generates phosphatidylinositol 3-phosphate (PIP3) and phosphatidylinositol 3,4-bisphosphate (PI(3, 4)P2) from phosphatidylinositol and phosphatidylinositol 4-phosphate (3). PI3K Class II  $\alpha$  is located in various intracellular locations such as the trans-Golgi network, endocytic compartments, clathrin-coated vesicles, and nuclear speckles (1,4,5). Research studies have indicated that PI3K Class II  $\alpha$  regulates the assembly and distribution of clathrin, resulting in the modulation of clathrin-dependent trafficking and sorting within the trans Golgi network (5,6). PI3K Class II  $\alpha$  also mediates translocation of the glucose transporter GLUT4 to the plasma membrane in response to insulin (7). PI3K Class II  $\alpha$  has also been shown to regulate neurosecretory granule exocytosis (8) and vascular smooth muscle contraction (9). Unlike other PI3K family members, PI3K Class II  $\alpha$  is less sensitive to the PI3K inhibitors wortmannin and LY294002 (3).

### Background References

- Didichenko, S.A. and Thelen, M. (2001) *J Biol Chem* 276, 48135-42.
- Stahelin, R.V. et al. (2006) *J Biol Chem* 281, 39396-406.
- Domin, J. et al. (1997) *Biochem J* 326 ( Pt 1), 139-47.
- Domin, J. et al. (2000) *J Biol Chem* 275, 11943-50.
- Gaidarov, I. et al. (2001) *Mol Cell* 7, 443-9.
- Gaidarov, I. et al. (2005) *J Biol Chem* 280, 40766-72.
- Falasca, M. et al. (2007) *J Biol Chem* 282, 28226-36.
- Wen, P.J. et al. (2008) *Mol Biol Cell* 19, 5593-603.
- Yoshioka, K. et al. (2007) *Mol Pharmacol* 71, 912-20.

### Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

### Western Blot Buffer

**IMPORTANT:** For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at  $4^{\circ}\text{C}$  with gentle shaking, overnight.

### Applications Key

**WB:** Western Blotting **IP:** Immunoprecipitation

### Cross-Reactivity Key

**H:** human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster  
**X:** Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse  
**GP:** Guinea Pig **Rab:** rabbit **All:** all species expected

### Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.  
All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](http://cellsignal.com/trademarks) for more information.

**Limited Uses**

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.