4/18/24, 10:34 AM Revision 3

-

e at -20C	MSH6 (3E1) Mouse mAb		Cell Signaling	
Store (		Orders:	877-616-CELL (2355) orders@cellsignal.com	
88		Support:	877-678-TECH (8324)	
#12988		Web:	info@cellsignal.com cellsignal.com	
#		3 Trask Lane   Danvers   Ma		

## For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: WB, IHC-P, IF-IC	Reactivity: H	Sensitivity: Endogenous	<b>MW (kDa):</b> 160	Source/Isotype: Mouse IgG2a	<b>UniProt ID:</b> #P52701	Entrez-Gene Id: 2956		
Product Usage	A	pplication				Dilution		
Information	W	estern Blotting				1:1000		
	In	nmunohistochemistry	(Paraffin)			1:500		
	In	nmunofluorescence (I	mmunocytochen		1:100			
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody. For a carrier free (BSA and azide free) version of this product see product #98142.						
	Fo							
Specificity / Sensitivity		MSH6 (3E1) Mouse mAb recognizes endogenous levels of total MSH6 protein.						
Source / Purification		Monoclonal antibody is produced by immunizing animals with a recombinant protein fragment specific to human MSH6 protein expressed in <i>E.coli</i> .						
Background		The DNA mismatch repair system (MMR) repairs post-replication DNA, inhibits recombination between nonidentical DNA sequences, and induces both checkpoint and apoptotic responses following certain types of DNA damage (1). MSH2 (MutS homologue 2) forms the hMutS-α dimer with MSH6 and is an essential component of the mismatch repair process. hMutS-α is part of the BRCA1-associated surveillance complex (BASC), a complex that also contains BRCA1, MLH1, ATM, BLM, PMS2 proteins, and the Rad50-Mre11-NBS1 complex (2). Mutations in MSH6 and other MMR proteins have been found in a large proportion of hereditary nonpolyposis colorectal cancer (Lynch Syndrome), the most common form of inherited colorectal cancer in the Western world (3). Mutations in MSH6 have been shown to occur in glioblastoma in response to temozolomide therapy and to promote temozolomide resistance (4).						
Background Refe	2. \ 3. I	O'Brien, V. and Browr Wang, Y. et al. (2000) Plotz, G. et al. (2006) Yip, S. et al. (2009) C	Genes Dev 14, J Mol Histol 37,	271-83.				
Species Reactivity		Species reactivity is determined by testing in at least one approved application (e.g., western blot).						
Western Blot Buffer		IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.						
Applications Key		WB: Western Blotting IHC-P: Immunohistochemistry (Paraffin) IF-IC: Immunofluorescence (Immunocytochemistry)						
Cross-Reactivity F	X:>	<ul> <li>H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster</li> <li>X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse</li> <li>GP: Guinea Pig Rab: rabbit All: all species expected</li> </ul>						
Trademarks and Patents	Sigr XP All c	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. SignalStain is a registered trademark of Cell Signaling Technology, Inc. XP is a registered trademark of Cell Signaling Technology, Inc. All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.						
Limited Uses								

## MSH6 (3E1) Mouse mAb (#12988) Datasheet Without Images Cell Signaling Technology

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.