#13162 store at -20C

S100A6 (D3H3W) Rabbit mAb



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: WB, IP	Reactivity: H M R Mk	Sensitivity: Endogenous	MW (kDa): 10	Source/Isotype: Rabbit IgG	UniProt ID: #P06703	Entrez-Gene Id: 6277	
Product Usage Information	Ар	plication		Dilution			
	We	stern Blotting		1:1000			
	Imr	nunoprecipitation		1:50			
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μ g/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20 °C. Do not aliquot the antibody.					
Specificity / Sensiti	- ,	S100A6 (D3H3W) Rabbit mAb recognizes endogenous levels of total S100A6 protein. This antibody also cross-reacts with unidentified proteins at 40 kDa and 60 kDa.					
Source / Purificatio		Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Asp65 of human S100A6 protein.					
Background		Despite their relatively small size (8-12 kDa) and uncomplicated architecture, S100 proteins regulate a variety of cellular processes, such as cell growth and motility, cell cycle progression, transcription, and					

Despite their relatively small size (8-12 kDa) and uncomplicated architecture, S100 proteins regulate a variety of cellular processes, such as cell growth and motility, cell cycle progression, transcription, and differentiation. To date, 25 members have been identified, including S100A1-S100A18, trichohyalin, filaggrin, repetin, S100P, and S100Z, making it the largest group in the EF-hand, calcium-binding protein family. Interestingly, 14 S100 genes are clustered on human chromosome 1q21, a region of genomic instability. Research studies have demonstrated that significant correlation exists between aberrant S100 protein expression and cancer progression. S100 proteins primarily mediate immune responses in various tissue types but are also involved in neuronal development (1-4).

Each S100 monomer bears two EF-hand motifs and can bind up to two molecules of calcium (or other divalent cation in some instances). Structural evidence shows that S100 proteins form antiparallel homo- or heterodimers that coordinate binding partner proximity in a calcium-dependent (and sometimes calcium-independent) manner. Although structurally and functionally similar, individual members show restricted tissue distribution, are localized in specific cellular compartments, and display unique protein binding partners, which suggests that each plays a specific role in various signaling pathways. In addition to an intracellular role, some S100 proteins have been shown to act as receptors for extracellular ligands or are secreted and exhibit cytokine-like activities (1-4).

S100A6 (calcyclin) is involved in a number of cellular processes, including exocytosis (5) and cell cycle regulation (6). In addition, S100A6 interacts with a number of proteins such as SIP, GAPDH, and annexins in a calcium-dependent fashion (7,8). Research studies demonstrate that a down regulation of corresponding S100A6 gene expression causes a decrease in cell proliferation (6).

Background References

- 1. Heizmann, C.W. et al. (2002) Front Biosci 7, d1356-68.
- 2. Donato, R. (2003) Microsc Res Tech 60, 540-51.
- 3. Marenholz, I. et al. (2004) Biochem Biophys Res Commun 322, 1111-22.
- 4. Santamaria-Kisiel, L. et al. (2006) Biochem J 396, 201-14.
- 5. Okazaki, K. et al. (1994) J Biol Chem 269, 6149-52.
- 6. Breen, E.C. and Tang, K. (2003) J Cell Biochem 88, 848-54.
- 7. Nowotny, M. et al. (2003) J Biol Chem 278, 26923-8.
- 8. Filipek, A. and Wojda, U. (1996) Biochem J 320 (Pt 2), 585-7.
- 9. Bao, L. et al. (2012) FEBS J 279, 4576-88.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

1/1/24. 12:56 PM

Western Blot Buffer

S100A6 (D3H3W) Rabbit mAb (#13162) Datasheet Without Images Cell Signaling Technology

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

Cross-Reactivity Key

WB: Western Blotting IP: Immunoprecipitation

H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster X: Xenopus Z: zebrafish B: bovine Dq: dog Pq: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse

GP: Guinea Pig Rab: rabbit All: all species expected

Trademarks and **Patents**

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

XP is a registered trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more

information.

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

> Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.

Limited Uses