

#13166 Store at -20°C

Integrin $\beta 3$ (D7X3P) XP® Rabbit mAb



Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
WB, IHC-P	H M	Endogenous	100	Rabbit IgG	#P05106	3690

Product Usage Information

Application

Western Blotting

Immunohistochemistry (Paraffin)

Dilution

1:1000

1:125 - 1:500

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μ g/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

For a carrier-free (BSA and azide free) version of this product see product #29462.

Specificity / Sensitivity

Integrin $\beta 3$ (D7X3P) XP® Rabbit mAb recognizes endogenous levels of total integrin $\beta 3$ protein.

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Ile114 of human integrin $\beta 3$ protein.

Background

Integrins are heterodimeric cell surface receptors that play a pivotal role in cell adhesion and migration, as well as in growth and survival (1,2). The integrin family contains at least 18 α and 8 β subunits that form 24 known integrins with distinct tissue distribution and overlapping ligand specificities (3). Integrins not only transmit signals to cells in response to the extracellular environment (outside-in signaling), but also sense intracellular cues to alter their interaction with the extracellular environment (inside-out signaling) (1,2). α IIb β 3 and α v β 3 are the two β 3 containing integrins which are prominently expressed in hematopoietic cells and angiogenic endothelial cells and perform adhesive functions in hemostasis, wound healing and angiogenesis (1,4). Tyr773 and Tyr785 (usually referred to as Tyr747 and Tyr759 based on the chicken sequence) are phosphorylated upon ligand binding (5). Phosphorylation of these tyrosine residues is required for certain ligand-induced signaling (6). Thr779 (corresponding to Thr753 of the chicken sequence) of integrin $\beta 3$ in the platelet specific α IIb β 3 is phosphorylated by PKD and/or Akt, which may modulate integrin association with other signaling molecules (7).

Background References

1. Liu, S. et al. (2000) *J Cell Sci* 113 (Pt 20), 3563-71.
2. Hood, J.D. and Chersesh, D.A. (2002) *Nat Rev Cancer* 2, 91-100.
3. van der Flier, A. and Sonnenberg, A. (2001) *Cell Tissue Res* 305, 285-98.
4. Shattil, S.J. (1995) *Thromb Haemost* 74, 149-55.
5. Blystone, S.D. (2002) *J Biol Chem* 277, 46886-90.
6. Butler, B. et al. (2003) *J Biol Chem* 278, 5264-70.
7. Kirk, R.I. et al. (2000) *J Biol Chem* 275, 30901-6.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting **IHC-P:** Immunohistochemistry (Paraffin)

Cross-Reactivity Key

H: human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster
X: Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse
GP: Guinea Pig **Rab:** rabbit **All:** all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.
XP is a registered trademark of Cell Signaling Technology, Inc.
All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.