

#13167 Store at -20°C

Phospho-GIT2 (Tyr592) (D6L1J) Rabbit mAb



Cell Signaling
TECHNOLOGY®

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: WB	Reactivity: H	Sensitivity: Endogenous	MW (kDa): 85	Source/Isotype: Rabbit IgG	UniProt ID: #Q14161	Entrez-Gene Id: 9815
---------------------	------------------	----------------------------	-----------------	-------------------------------	------------------------	-------------------------

Product Usage Information	Application Western Blotting	Dilution 1:1000
Storage	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.	
Specificity / Sensitivity	Phospho-GIT2 (Tyr592) (D6L1J) Rabbit mAb recognizes endogenous levels of GIT2 protein only when phosphorylated at Tyr592.	
Source / Purification	Monoclonal antibody is produced by immunizing animals with a synthetic phosphopeptide corresponding to residues surrounding Tyr592 of human GIT2 protein.	
Background	G protein-coupled receptor (GPCR) kinase interacting proteins 1 and 2 (GIT1 and GIT2) are highly conserved, ubiquitous scaffold proteins involved in localized signaling to help regulate focal contact assembly and cytoskeletal dynamics. GIT proteins contain multiple interaction domains that allow interaction with small GTPases (including ARF, Rac, and cdc42), kinases (such as PAK and MEK), the Rho family GEF Pix, and the focal adhesion protein paxillin (reviewed in 1). GIT1 and GIT2 share many of the same properties, but with at least ten distinct, tissue-specific splice variants. GIT2 has been shown to play an important role inhibiting focal adhesion turnover and membrane protrusion (2,3). Focal adhesion localization and paxillin binding of GIT2 is regulated through phosphorylation at one or more tyrosine sites (Tyr286, Tyr392, Tyr592) by FAK and/or Src (4,5, reviewed in 6). Once at the focal adhesion, GIT2 is thought to play a key role in cell polarity and migration, making it a protein of interest in the investigation of oncogenic signaling pathways (3,5,7).	
Background References	<ol style="list-style-type: none"> 1. Hoefen, R.J. and Berk, B.C. (2006) <i>J Cell Sci</i> 119, 1469-75. 2. Premont, R.T. et al. (2000) <i>J Biol Chem</i> 275, 22373-80. 3. Frank, S.R. et al. (2006) <i>EMBO J</i> 25, 1848-59. 4. Brown, M.C. et al. (2005) <i>Mol Biol Cell</i> 16, 4316-28. 5. Yu, J.A. et al. (2009) <i>Mol Biol Cell</i> 20, 4706-19. 6. Yu, J.A. et al. (2010) <i>Cell Adh Migr</i> 4, 342-7. 7. Mazaki, Y. et al. (2006) <i>Nat Immunol</i> 7, 724-31. 	

Species Reactivity	Species reactivity is determined by testing in at least one approved application (e.g., western blot).
Western Blot Buffer	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.
Applications Key	WB: Western Blotting
Cross-Reactivity Key	H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse GP: Guinea Pig Rab: rabbit All: all species expected
Trademarks and Patents	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.
Limited Uses	Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.