

#13492 Store at -20C

Atg101 (E1Z4W) Rabbit mAb



Cell Signaling
TECHNOLOGY®

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
WB	H M R Mk	Endogenous	25	Rabbit IgG	#Q9BSB4	60673

Product Usage Information

Application

Western Blotting

Dilution

1:1000

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Specificity / Sensitivity

Atg101 (E1Z4W) Rabbit mAb recognizes endogenous levels of total Atg101 protein.

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Val177 of human Atg101 protein.

Background

Atg101 was discovered as a binding protein for Atg13, a component of the ULK1 serine-threonine kinase complex required for autophagy (1-3). Autophagy is a catabolic process for the autophagosomic-lysosomal degradation of bulk cytoplasmic contents (4,5). It is generally activated by conditions of nutrient deprivation, but is also associated with a number of physiological processes, including development, differentiation, neurodegeneration, infection, and cancer (6). The molecular machinery of autophagy was largely discovered in yeast and is directed by a number of autophagy-related (Atg) genes. The ULK1 complex includes both Atg13 and FIP200 and is required for starvation-induced autophagy (7-9). Interaction between Atg101 and Atg13 can be important for the stability and basal phosphorylation of Atg13 and ULK1 (1,2).

Background References

1. Mercer, C.A. et al. (2009) *Autophagy* 5, 649-62.
2. Hosokawa, N. et al. (2009) *Autophagy* 5, 973-9.
3. Chan, E.Y. et al. (2007) *J Biol Chem* 282, 25464-74.
4. Reggiori, F. and Klionsky, D.J. (2002) *Eukaryot Cell* 1, 11-21.
5. Codogno, P. and Meijer, A.J. (2005) *Cell Death Differ* 12 Suppl 2, 1509-18.
6. Levine, B. and Yuan, J. (2005) *J Clin Invest* 115, 2679-88.
7. Hosokawa, N. et al. (2009) *Mol Biol Cell* 20, 1981-91.
8. Jung, C.H. et al. (2009) *Mol Biol Cell* 20, 1992-2003.
9. Ganley, I.G. et al. (2009) *J Biol Chem* 284, 12297-305.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting

Cross-Reactivity Key

H: human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster
X: Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse
GP: Guinea Pig **Rab:** rabbit **All:** all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.
All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.