

#13531 Store at -20°C

Afadin (D1Y3Z) Rabbit mAb**Cell Signaling**
TECHNOLOGY®**Orders:** 877-616-CELL (2355)
orders@cellsignal.com**Support:** 877-678-TECH (8324)**Web:** info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
WB, IP	H M R Mk Dg	Endogenous	205	Rabbit IgG	#P55196	4301

Product Usage Information**Application**Western Blotting
Immunoprecipitation**Dilution**1:1000
1:50**Storage**

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Specificity / Sensitivity

Afadin (D1Y3Z) Rabbit mAb recognizes endogenous levels of total afadin protein. Based on the protein sequence, this antibody is expected to recognize all afadin isoforms.

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Arg1117 of human afadin protein.

Background

In multicellular organisms, intercellular junctions play essential roles in tissue integrity and maintenance of cell polarity. Tight junctions (TJs) form a continuous barrier to fluids across the epithelium and endothelium (reviewed in 1). Adherens junctions (AJs) are dynamic structures that form cell-cell contacts linking cells into a continuous sheet (reviewed in 2). The actin filament-binding protein, Afadin, binds to nectin forming a connection to the actin cytoskeleton (3). AJs are formed when nectin assembles cadherin at the cell-cell adhesion site and these junctions are then involved in the formation and maintenance of TJs (4,5). Afadin has two splice variants: l-afadin, which is ubiquitously expressed, and s-afadin, which is expressed predominantly in neural tissue. s-Afadin is a shorter form lacking one of the three proline-rich regions found in l-afadin, as well as the carboxyl-terminal F-actin binding region (6). Human s-afadin is identical to AF-6, the ALL-1 fusion partner involved in acute myeloid leukemias (7). Recent work has also shown that afadin is involved in controlling the directionality of cell movement when it is localized at the leading edge of moving cells (8,9).

Background References

- Shin, K. et al. (2006) *Annu Rev Cell Dev Biol* 22, 207-35.
- Harris, T.J. and Tepass, U. (2010) *Nat Rev Mol Cell Biol* 11, 502-14.
- Ikeda, W. et al. (1999) *J Cell Biol* 146, 1117-32.
- Sato, T. et al. (2006) *J Biol Chem* 281, 5288-99.
- Ooshio, T. et al. (2007) *J Cell Sci* 120, 2352-65.
- Mandai, K. et al. (1997) *J Cell Biol* 139, 517-28.
- Prasad, R. et al. (1993) *Cancer Res* 53, 5624-8.
- Miyata, M. et al. (2009) *J Cell Sci* 122, 4319-29.
- Miyata, M. et al. (2009) *J Biol Chem* 284, 24595-609.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key**WB:** Western Blotting **IP:** Immunoprecipitation**Cross-Reactivity Key**

H: human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster
X: Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse
GP: Guinea Pig **Rab:** rabbit **All:** all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.
 XP is a registered trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.