#13567 store at +4c

Phospho-RelB (Ser552) (D41B9) XP[®] Rabbit mAb (PE Conjugate)



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: Reactivity: Sensitivity: Source/Isotype: **UniProt ID:** Entrez-Gene Id: FC-FP HMEndogenous Rabbit IgG #Q01201 5971 **Product Usage** Application Dilution Information Flow Cytometry (Fixed/Permeabilized) 1:50 Supplied in PBS (pH 7.2), less than 0.1% sodium azide and 2 mg/ml BSA. Store at 4°C. Do not aliquot the **Storage** antibodies. Protect from light. Do not freeze.

Specificity / Sensitivity Phospho-RelB (Ser552) (D41B9) XP® Rabbit mAb (PE Conjugate) detects endogenous levels of RelB

protein only when phosphorylated at Ser552.

Species predicted to react based on 100% sequence homology:

Rat, Monkey, Bovine, Dog

Source / Purification Monoclonal antibody is produced by immunizing animals with a synthetic phosphopeptide corresponding to

residues surrounding Ser552 of mouse RelB protein.

Product DescriptionThis Cell Signaling Technology antibody is conjugated to phycoerythrin (PE) and tested in-house for direct flow cytometry analysis in human cells. The antibody is expected to exhibit the same species cross-

reactivity as the unconjugated Phospho-RelB (Ser552) (D41B9) XP® Rabbit mAb #5025.

Background Transcription factors of the nuclear factor κB (NF-κB)/Rel family play a pivotal role in inflammatory and

immune responses (1,2). There are five family members in mammals: ReIA, c-ReI, ReIB, NF-κB1 (p105/p50), and NF-κB2 (p100/p52). Both p105 and p100 are proteolytically processed by the proteasome to produce p50 and p52, respectively. Rel proteins bind p50 and p52 to form dimeric complexes that bind DNA and regulate transcription. In unstimulated cells, NF-κB is sequestered in the cytoplasm by IκB inhibitory proteins (3-5). NF-κB-activating agents can induce the phosphorylation of IκB proteins, targeting them for rapid degradation through the ubiquitin-proteasome pathway and releasing NF-κB to enter the nucleus where it regulates gene expression (6-8). NIK and IKKα (IKK1) regulate the phosphorylation and

processing of NF-kB2 (p100) to produce p52, which translocates to the nucleus (9-11).

Background References 1. Baeuerle, P.A. and Henkel, T. (1994) Annu Rev Immunol 12, 141-79.

2. Baeuerle, P.A. and Baltimore, D. (1996) Cell 87, 13-20.

3. Haskill, S. et al. (1991) Cell 65, 1281-9.

4. Thompson, J.E. et al. (1995) Cell 80, 573-82.

5. Whiteside, S.T. et al. (1997) *EMBO J* 16, 1413-26.

6. Traenckner, E.B. et al. (1995) EMBO J 14, 2876-83.

7. Scherer, D.C. et al. (1995) Proc Natl Acad Sci USA 92, 11259-63.

8. Chen, Z.J. et al. (1996) Cell 84, 853-62.

9. Senftleben, U. et al. (2001) Science 293, 1495-9.

10. Coope, H.J. et al. (2002) EMBO J 21, 5375-85.

11. Xiao, G. et al. (2001) Mol Cell 7, 401-9.

Species Reactivity Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Applications Key FC-FP: Flow Cytometry (Fixed/Permeabilized)

Cross-Reactivity Key

H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse GP: Guinea Pig Rab: rabbit All: all species expected

Trademarks and Patents

Limited Uses

 $\label{lem:cell_signaling} \textbf{ Technology is a trademark of Cell Signaling Technology, Inc.}$

XP is a registered trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.