

#13768 Store at -20°C

**PHC1 (1F3F3) Mouse mAb****Cell Signaling**  
TECHNOLOGY®**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com**Support:** 877-678-TECH (8324)**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

**For Research Use Only. Not for Use in Diagnostic Procedures.**

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
WB, IP, ChIP, ChIP-seq	H M	Endogenous	130	Mouse IgG2b	#P78364	1911

**Product Usage Information**

For optimal ChIP and ChIP-seq results, use 10 µl of antibody and 10 µg of chromatin (approximately 4 x 10<sup>6</sup> cells) per IP. This antibody has been validated using SimpleChIP® Enzymatic Chromatin IP Kits.

Application	Dilution
Western Blotting	1:1000
Immunoprecipitation	1:50
Chromatin IP	1:50
Chromatin IP-seq	1:50

**Storage**

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. *Do not aliquot the antibody.*

**Specificity / Sensitivity**

PHC1 (1F3F3) Mouse mAb recognizes endogenous levels of total PHC1 protein.

**Source / Purification**

Monoclonal antibody is produced by immunizing animals with recombinant protein specific to the carboxy terminus of human PHC1 protein.

**Background**

The polycomb group (PcG) proteins contribute to the maintenance of cell identity, stem cell self-renewal, cell cycle regulation, and oncogenesis by maintaining the silenced state of genes that promote cell lineage specification, cell death, and cell-cycle arrest (1-4). Polycomb group proteins regulate cell proliferation and senescence through repression of the p16Ink4a and p19Arf genes, and are essential in maintaining adult hematopoietic, neural stem cells, and embryonic stem cells (3-5). PcG proteins are found in two complexes that cooperate to maintain long-term gene silencing through epigenetic chromatin modifications. DNA-binding transcription factors recruit the EED-EZH2 complex to genes, which methylates histone H3 on Lys27 (6). Methylation of Lys27 facilitates the recruitment of the PRC1 complex, which ubiquitinylates histone H2A on Lys119 (7). PRC1 is composed of BMI1 and RING1A, which enhance the E3 ubiquitin ligase activity of the RING1B catalytic subunit (8). Polyhomeotic-like 1 (PHC1) is one of several additional PRC1 complex proteins that are required to maintain the silenced state of PRC1 target genes and mediate proper anterior-posterior specification during development (9). Mutations in the corresponding *PHC1* gene correlate with an autosomal recessive form of primary microcephaly characterized by low-to-normal cognitive function and impaired DNA repair (10).

**Background References**

1. Boyer, L.A. et al. (2006) *Nature* 441, 349-53.
2. Lee, T.I. et al. (2006) *Cell* 125, 301-13.
3. Park, I.K. et al. (2003) *Nature* 423, 302-5.
4. Molofsky, A.V. et al. (2003) *Nature* 425, 962-7.
5. Molofsky, A.V. et al. (2005) *Genes Dev* 19, 1432-7.
6. Cao, R. and Zhang, Y. (2004) *Mol Cell* 15, 57-67.
7. Wang, H. et al. (2004) *Nature* 431, 873-8.
8. Cao, R. et al. (2005) *Mol Cell* 20, 845-54.
9. Isono, K. et al. (2005) *Mol Cell Biol* 25, 6694-706.
10. Awad, S. et al. (2013) *Hum Mol Genet* 22, 2200-13.

**Species Reactivity**

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

**Western Blot Buffer**

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

**Applications Key**

**WB:** Western Blotting **IP:** Immunoprecipitation **ChIP:** Chromatin IP **ChIP-seq:** Chromatin IP-seq

**Cross-Reactivity Key**

**H:** human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster  
**X:** Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse  
**GP:** Guinea Pig **Rab:** rabbit **All:** all species expected

**Trademarks and Patents**

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

SimpleChIP is a registered trademark of Cell Signaling Technology, Inc.

XP is a registered trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](https://cellsignal.com/trademarks) for more information.

**Limited Uses**

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.