

**#13826** Store at -20°C

## ERRα (E1G1J) Rabbit mAb


**Cell Signaling**  
TECHNOLOGY®

**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com

**Support:** 877-678-TECH (8324)

**Web:** info@cellsignal.com  
cellsignal.com

**3 Trask Lane | Danvers | Massachusetts | 01923 | USA**
**For Research Use Only. Not for Use in Diagnostic Procedures.**

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
WB, ChIP, ChIP-seq	H M R Mk	Endogenous	50	Rabbit IgG	#P11474	2101

### Product Usage Information

For optimal ChIP and ChIP-seq results, use 10 µl of antibody and 10 µg of chromatin (approximately 4 x 10<sup>6</sup> cells) per IP. This antibody has been validated using SimpleChIP® Enzymatic Chromatin IP Kits.

Application	Dilution
Western Blotting	1:1000
Chromatin IP	1:50
Chromatin IP-seq	1:50

### Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

### Specificity / Sensitivity

ERRα (E1G1J) Rabbit mAb recognizes endogenous levels of total ERRα protein. This antibody does not cross-react with ERR family members ERRβ and ERRγ, and does not cross-react with either ERα or ERβ.

### Species predicted to react based on 100% sequence homology:

Bovine, Dog, Pig, Horse

### Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues near the carboxy terminus of human ERRα protein.

### Background

The estrogen-related receptor (ERR) subfamily of orphan nuclear receptors include three protein receptors, ERRα/NR3B1, ERRβ/NR3B2, and ERRγ/NR3B3, that have yet to be associated with natural ligands. PGC-1 coactivators regulate ERR transcription activation ability and receptor-induced transcription of genes involved in lipid metabolism, glucose metabolism, and mitochondrial biogenesis (1). Estrogen-related receptor α (ERRα/NR3B1) is an orphan nuclear receptor that controls transcription of genes involved in fatty acid oxidation, glucose metabolism, and mitochondrial biogenesis (1,2). The receptor protein contains a non-conserved amino terminal domain (NTD), a central zinc finger DNA binding domain, and a ligand-binding domain. The carboxy-terminal AF2 helix motif of ERRα contains binding sites for nuclear receptor coactivators PGC-1α and PGC-1β (3-5). Research studies demonstrate that ERRα transcriptional activity is regulated through phosphorylation and sumoylation within the NTD (6). ERRα is ubiquitously expressed, with strong expression observed in heart, kidneys, skeletal muscle, and other high metabolic demand tissues (2). Additional studies indicate that ERRα is coexpressed in breast tumors with unfavorable biomarkers (7). The pharmacologic inhibition of ERRα activity in breast cancer might serve as a valuable therapeutic approach (8,9).

### Background References

1. Giguère, V. (2008) *Endocr Rev* 29, 677-96.
2. Giguère, V. et al. (1988) *Nature* 331, 91-4.
3. Huss, J.M. et al. (2002) *J Biol Chem* 277, 40265-74.
4. Schreiber, S.N. et al. (2003) *J Biol Chem* 278, 9013-8.
5. Kamei, Y. et al. (2003) *Proc Natl Acad Sci U S A* 100, 12378-83.
6. Tremblay, A.M. et al. (2008) *Mol Endocrinol* 22, 570-84.
7. Ariazi, E.A. et al. (2002) *Cancer Res* 62, 6510-8.
8. Chang, C.Y. et al. (2011) *Cancer Cell* 20, 500-10.
9. Deblois, G. et al. (2009) *Cancer Res* 69, 6149-57.
10. Lanvin, O. et al. (2007) *J Biol Chem* 282, 28328-34.
11. Willy, P.J. et al. (2004) *Proc Natl Acad Sci U S A* 101, 8912-7.

### Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

**Western Blot Buffer**

**IMPORTANT:** For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

**Applications Key**

**WB:** Western Blotting **ChIP:** Chromatin IP **ChIP-seq:** Chromatin IP-seq

**Cross-Reactivity Key**

**H:** human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster  
**X:** Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse  
**GP:** Guinea Pig **Rab:** rabbit **All:** all species expected

**Trademarks and Patents**

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

SimpleChIP is a registered trademark of Cell Signaling Technology, Inc.

XP is a registered trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](https://cellsignal.com/trademarks) for more information.

**Limited Uses**

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.