

#13808 Store at -20°C

**PTPN14 (D5T6Y) Rabbit mAb****Cell Signaling**  
TECHNOLOGY®**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com**Support:** 877-678-TECH (8324)**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

**For Research Use Only. Not for Use in Diagnostic Procedures.**

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
WB	H M R	Endogenous	160	Rabbit IgG	#Q15678	5784

**Product Usage Information****Application**

Western Blotting

**Dilution**

1:1000

**Storage**Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. *Do not aliquot the antibody.***Specificity / Sensitivity**

PTPN14 (D5T6Y) Rabbit mAb recognizes endogenous levels of total PTPN14 protein.

**Source / Purification**

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Leu737 of human PTPN14 protein.

**Background**

Tyrosine-protein phosphatase non-receptor type-14 (PTPN14, Pez, PTPD2 and PTP36) is an evolutionarily conserved non-membrane tyrosine phosphatase with homology to the band 4.1 family of proteins (1-3). The PTPN14 protein contains an amino-terminal FERM (4.1-ezrin-radixin-moesin) domain, which suggests plasma membrane localization of the protein, and a carboxy-terminal protein tyrosine phosphatase (PTP) domain (4). Research studies have identified possible roles for PTPN14 in multiple, diverse signaling pathways, including cell growth and proliferation, cell migration and adhesion, and development. The PTPN14 phosphatase regulates the subcellular localization of YAP in a cell density-dependent manner, indicating a role for PTPN14 in the Hippo signaling pathway (5). The *Drosophila* PTPN14 homolog Pez localizes to adherens junctions, where it may regulate cell motility through dephosphorylation of  $\beta$ -catenin (3). PTPN14 may play a role in epithelial-mesenchymal transition through effects on the TGF- $\beta$  signaling pathway (6), and interacts with VEGFR3, a receptor tyrosine kinase involved in lymphangiogenesis (7). Loss-of-function mutations in the *PTPN14* gene are associated with colorectal cancer (8), and choanal atresia and lymphedema, an autosomal recessive disorder characterized by defects in both nasal passage development and lymphangiogenesis (7).

**Background References**

- Smith, A.L. et al. (1995) *Biochem Biophys Res Commun* 209, 959-65.
- Ogata, M. et al. (1999) *J Biol Chem* 274, 20717-24.
- Wadham, C. et al. (2003) *Mol Biol Cell* 14, 2520-9.
- Wadham, C. et al. (2000) *J Cell Sci* 113 ( Pt 17), 3117-23.
- Wang, W. et al. (2012) *Genes Dev* 26, 1959-71.
- Wyatt, L. et al. (2007) *J Cell Biol* 178, 1223-35.
- Au, A.C. et al. (2010) *Am J Hum Genet* 87, 436-44.
- Laczmanska, I. and Sasiadek, M.M. (2011) *Acta Biochim Pol* 58, 467-70.

**Species Reactivity**

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

**Western Blot Buffer****IMPORTANT:** For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.**Applications Key****WB:** Western Blotting**Cross-Reactivity Key**

**H:** human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster  
**X:** Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse  
**GP:** Guinea Pig **Rab:** rabbit **All:** all species expected

**Trademarks and Patents**

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.  
 All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](http://cellsignal.com/trademarks) for more information.

**Limited Uses**

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.