

#13945 Store at -20°C

INTS9 Antibody



Cell Signaling
TECHNOLOGY®

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: WB, IP, ChIP	Reactivity: H M R Mk	Sensitivity: Endogenous	MW (kDa): 75	Source: Rabbit	UniProt ID: #Q9NV88	Entrez-Gene Id: 55756
--------------------------------------	--------------------------------	-----------------------------------	------------------------	--------------------------	-------------------------------	---------------------------------

Product Usage Information

For optimal ChIP results, use 10 µl of antibody and 10 µg of chromatin (approximately 4 x 10⁶ cells) per IP. This antibody has been validated using SimpleChIP® Enzymatic Chromatin IP Kits.

Application	Dilution
Western Blotting	1:1000
Immunoprecipitation	1:100
Chromatin IP	1:50

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at –20°C. Do not aliquot the antibody.

Specificity / Sensitivity

INTS9 Antibody recognizes endogenous levels of total INTS9 protein.

Species predicted to react based on 100% sequence homology:

Hamster, Bovine, Dog, Pig

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues surrounding His601 of human INTS9 protein. Antibodies are purified by protein A and peptide affinity chromatography.

Background

The integrator complex is an evolutionarily conserved complex that is composed of at least 12 subunits in humans. It is thought to be a multifunctional complex with roles in orchestrating snRNA 3' end processing with transcription termination, DNA double-stranded break repair, hematopoietic development, and cell cycle progression (1-6). The integrator subunits (INTS) 9 and 11 are thought to be the catalytic subunits of the complex and are essential for the function of the complex (6,7). Research studies indicate that the integrator complex is recruited to snRNA genes through its interaction with the carboxy-terminal domain (CTD) of Rpb1, the largest subunit of RNA polymerase II (8). Phosphorylation of the Rpb1 CTD heptapeptide repeat residues Ser2 and Ser7 is required for efficient binding of integrator subunit proteins (9).

Background References

- Chen, J. and Wagner, E.J. (2010) *Biochem Soc Trans* 38, 1082-7.
- O'Reilly, D. et al. (2014) *Nucleic Acids Res* 42, 264-75.
- Tao, S. et al. (2009) *Development* 136, 2757-65.
- Huang, J. et al. (2009) *Mol Cell* 35, 384-93.
- Li, Y. et al. (2009) *J Biol Chem* 284, 23525-31.
- Dominski, Z. et al. (2005) *Mol Cell Biol* 25, 1489-500.
- Albrecht, T.R. and Wagner, E.J. (2012) *Mol Cell Biol* 32, 1112-23.
- Baillat, D. et al. (2005) *Cell* 123, 265-76.
- Egloff, S. et al. (2010) *J Biol Chem* 285, 20564-9.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting **IP:** Immunoprecipitation **ChIP:** Chromatin IP

Cross-Reactivity Key

H: human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster
X: Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse
GP: Guinea Pig **Rab:** rabbit **All:** all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.
SimpleChIP is a registered trademark of Cell Signaling Technology, Inc.
All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.