PSMD11 (D1T1R) Rabbit mAb



Orders:

877-616-CELL (2355) orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

Applications: WB	Reactivity: H M R Mk	Sensitivity: Endogenous	MW (kDa): 47	Source/Isotype: Rabbit IgG	UniProt ID: #O00231	Entrez-Gene Id 5717	
Product Usage	Ар	plication			Dilution		
Information	We	estern Blotting		1:1000			
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μ g/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20° C. Do not aliquot the antibody.					
Specificity / Sensitivity		PSMD11 (D1T1R) Rabbit mAb recognizes endogenous levels of total PSMD11 protein. This antibody does not cross-react with COPS2 protein.					
Species predicted react based on 100 sequence homolog)%	rine, Horse					
Source / Purification		,	,	munizing animals with a synthetic peptide corresponding to man PSMD11 protein.			
Background	The	The 26S proteasome is a highly abundant proteolytic complex involved in the degradation of ubiquitinated					

substrate proteins. It consists largely of two sub-complexes, the 20S catalytic core particle (CP) and the 19S/PA700 regulatory particle (RP) that can cap either end of the CP. The CP consists of two stacked heteroheptameric β -rings (β_{1-7}) that contain three catalytic β -subunits and are flanked on either side by two heteroheptameric α -rings (α_{1-7}). The RP includes a base and a lid, each having multiple subunits. The base, in part, is composed of a heterohexameric ring of ATPase subunits belonging to the AAA (ATPases Associated with diverse cellular Activities) family. The ATPase subunits function to unfold the substrate and open the gate formed by the α-subunits, thus exposing the unfolded substrate to the catalytic β-subunits. The lid consists of ubiquitin receptors and DUBs that function in recruitment of ubiquitinated substrates and modification of ubiquitin chain topology (1,2). Other modulators of proteasome activity, such as PA28/11S

REG, can also bind to the end of the 20S CP and activate it (1,2).

Proteasome 26S subunit, non-ATPase, 11 (PSMD11, RPN6) is a non-ATPase subunit of the 19S/PA700 RP lid subcomplex. Research studies have shown that PSMD11 expression is regulated by FOXO4 in human embryonic stem cells (hESCs). Increased PSMD11 expression in hESCs correlates with high levels of proteasome activity, which may be required for protection from proteotoxic stress and senescence (3).

Background References

- 1. Finley, D. (2009) Annu Rev Biochem 78, 477-513.
- 2. Lee, M.J. et al. (2011) Mol Cell Proteomics 10, R110.003871.
- 3. Vilchez, D. et al. (2012) Nature 489, 304-8.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting

Cross-Reactivity Key

H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse

GP: Guinea Pig Rab: rabbit All: all species expected

Trademarks and **Patents**

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

PSMD11 (D1T1R) Rabbit mAb (#14303) Datasheet Without Images Cell Signaling Technology

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.