

Store at  
-20°C**PTMScan® IAP Buffer (10X)**Cell Signaling  
TECHNOLOGY®

#9993

0.6 mL

**Support:** +1-978-867-2388 (U.S.)  
[www.cellsignal.com/support](http://www.cellsignal.com/support)**Orders:** 877-616-2355 (U.S.)  
[orders@cellsignal.com](mailto:orders@cellsignal.com)**For Research Use Only. Not For Use In Diagnostic Procedures.****Description:** PTMScan® IAP Buffer is used to reconstitute lyophilized peptides prior to immunoaffinity purification (IAP).**Directions for Use:** Thaw 10X PTMScan® IAP buffer at room temperature or at 37 °C. Ensure all contents are dissolved as some components may have precipitated upon freezing. If necessary, dissolve by shaking gently. Before use, dilute with Milli-Q or equivalently purified H<sub>2</sub>O to 1X buffer (e.g., pipet 0.5 mL 10X IAP buffer from the vial into 4.5 mL H<sub>2</sub>O).**Solutions and Reagents:** 1X Buffer Components:

50 mM MOPS/NaOH pH 7.2

10 mM Na<sub>2</sub>HPO<sub>4</sub>

50 mM NaCl

**Storage:** This product is stable for 60 months when stored at -20°C. Prepared 1X IAP Buffer can be stored up to one month at 4°C.Thank you for your recent purchase. If you would like to provide a review visit [www.cellsignal.com/comments](http://www.cellsignal.com/comments).[www.cellsignal.com](http://www.cellsignal.com)