

#14457 Store at -20°C

ELMO1 (D4K2E) Rabbit mAb**Cell Signaling**
TECHNOLOGY®**Orders:** 877-616-CELL (2355)
orders@cellsignal.com**Support:** 877-678-TECH (8324)**Web:** info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
WB, IP	H	Endogenous	80	Rabbit IgG	#Q92556	9844

Product Usage Information**Application**Western Blotting
Immunoprecipitation**Dilution**1:1000
1:50**Storage**Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. *Do not aliquot the antibody.***Specificity / Sensitivity**

ELMO1 (D4K2E) Rabbit mAb recognizes endogenous levels of total ELMO1 protein.

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Ala567 of human ELMO1 protein.

Background

Engulfment and cell motility 1 (ELMO1) is a cell motility and migration protein that interacts with DOCK180 to form an atypical, two-part guanine nucleotide exchange factor (GEF) for the small GTPase Rac (1). The resultant localized Rac activation allows actin nucleation via WAVE family proteins, signaling to integrins, formation of lamellipodia and filopodia, and regulation of processes including phagocytosis and cell migration (2-4). Research studies indicate that DOCK180 and ELMO1 regulate cell migration in lymphocytes (5) and in ovarian cancer cells (6). ELMO1 also promotes Rac1-dependent cell motility through its interaction with the adaptor protein Nck-1 (7), and binds Arhgef16 to promote RhoG/Rac1-dependent engulfment of apoptotic cells by phagocytes (8). Polymorphisms in the corresponding *ELMO1* gene may be associated with susceptibility to diabetic neuropathy seen in selected populations of type II diabetic patients (9,10).

Background References

1. Brugnera, E. et al. (2002) *Nat Cell Biol* 4, 574-82.
2. Takenawa, T. and Miki, H. (2001) *J Cell Sci* 114, 1801-9.
3. Albert, M.L. et al. (2000) *Nat Cell Biol* 2, 899-905.
4. Gustavsson, A. et al. (2004) *J Biol Chem* 279, 22893-901.
5. Stevenson, C. et al. (2014) *J Immunol* 192, 6062-70.
6. Wang, J. et al. (2014) *Int J Gynecol Cancer* 24, 844-50.
7. Zhang, G. et al. (2014) *J Biol Chem* 289, 23112-22.
8. Lee, J. et al. (2014) *Biochim Biophys Acta* 1843, 2438-2447.
9. Pezzolesi, M.G. et al. (2009) *Diabetes* 58, 2698-702.
10. Wu, H.Y. et al. (2013) *J Endocrinol Invest* 36, 298-302.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key**WB:** Western Blotting **IP:** Immunoprecipitation**Cross-Reactivity Key**

H: human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster
X: Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse
GP: Guinea Pig **Rab:** rabbit **All:** all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.
 XP is a registered trademark of Cell Signaling Technology, Inc.
 All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.