

#13650 Store at -20C

# Langerin (D9H7R) XP® Rabbit mAb



**Cell Signaling**  
TECHNOLOGY®

**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com

**Support:** 877-678-TECH (8324)

**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

**For Research Use Only. Not for Use in Diagnostic Procedures.**

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
WB, IP, IF-IC, FC-FP	H	Endogenous	40-50	Rabbit IgG	#Q9UJ71	50489

## Product Usage Information

### Application

Western Blotting  
Immunoprecipitation  
Immunofluorescence (Immunocytochemistry)  
Flow Cytometry (Fixed/Permeabilized)

### Dilution

1:1000  
1:100  
1:800  
1:3200

## Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

For a carrier free (BSA and azide free) version of this product see product #49830.

## Specificity / Sensitivity

Langerin (D9H7R) XP® Rabbit mAb recognizes endogenous levels of total Langerin protein.

## Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Val14 of human Langerin protein.

## Background

Langerin (CD207) is a C-type lectin receptor whose expression is restricted mainly to dendritic cells in the skin including Langerhans cells in the epidermis and langerin+/CD103+ dermal dendritic cells (1-4). Langerin is found on the cell surface and within rod-shaped organelles called Birbeck granules, and its expression is required for the formation of Birbeck granules (1,5). Langerin recognizes carbohydrate motifs on the surface of pathogens, resulting in endocytosis of the pathogen into Birbeck granules, degradation of the pathogen, and antigen presentation to T cells (6-8).

## Background References

- Valladeau, J. et al. (2000) *Immunity* 12, 71-81.
- Poulin, L.F. et al. (2007) *J Exp Med* 204, 3119-31.
- Ginhoux, F. et al. (2007) *J Exp Med* 204, 3133-46.
- Bursch, L.S. et al. (2007) *J Exp Med* 204, 3147-56.
- Kissenpfennig, A. et al. (2005) *Mol Cell Biol* 25, 88-99.
- Stambach, N.S. and Taylor, M.E. (2003) *Glycobiology* 13, 401-10.
- Idoyaga, J. et al. (2008) *J Immunol* 180, 3647-50.
- Igyártó, B.Z. et al. (2011) *Immunity* 35, 260-72.

## Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

## Western Blot Buffer

**IMPORTANT:** For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

## Applications Key

**WB:** Western Blotting **IP:** Immunoprecipitation **IF-IC:** Immunofluorescence (Immunocytochemistry)  
**FC-FP:** Flow Cytometry (Fixed/Permeabilized)

## Cross-Reactivity Key

**H:** human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster  
**X:** Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse  
**GP:** Guinea Pig **Rab:** rabbit **All:** all species expected

## Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.  
Alexa Fluor is a registered trademark of Life Technologies Corporation.  
All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](http://cellsignal.com/trademarks) for more information.

## Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.