#14579 Store at +4C

Background

CD45 (Intracellular Domain) (D9M8I) XP[®] Rabbit mAb (PE Conjugate)



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: Reactivity: Sensitivity: Source/Isotype: **UniProt ID:** Entrez-Gene Id: FC-FP H Mk Endogenous Rabbit IgG #P08575 5788 **Product Usage** Application Dilution Information Flow Cytometry (Fixed/Permeabilized) 1:50

Storage Supplied in PBS (pH 7.2), less than 0.1% sodium azide and 2 mg/ml BSA. Store at 4°C. Do not aliquot the antibodies. Protect from light. Do not freeze.

Specificity / Sensitivity CD45 (Intracellular Domain) (D9M8I) XP® Rabbit mAb (PE Conjugate) recognizes endogenous levels of total CD45 protein.

Source / PurificationMonoclonal antibody is produced by immunizing animals with recombinant protein specific to the carboxy terminus of human CD45 protein.

Product Description

This Cell Signaling Technology antibody is conjugated to phycoerythrin (PE) and tested in-house for direct flow cytometry analysis in human cells. The antibody is expected to exhibit the same species cross-

reactivity as the unconjugated CD45 (Intracellular Domain) (D9M8I) XP^{\otimes} Rabbit mAb #13917.

The protein phosphatase (PTP) receptor CD45 is a type I transmembrane protein comprised of a pair of intracellular tyrosine phosphatase domains and a variable extracellular domain generated by alternative splicing (1). The catalytic activity of CD45 is a function of the first phosphatase domain (D1) while the second phosphatase domain (D2) may interact with and stabilize the first domain, or recruit/bind substrates (2,3). CD45 interacts directly with antigen receptor complex proteins or activates Src family kinases involved in the regulation of T- and B-cell antigen receptor signaling (1). Specifically, CD45 dephosphorylates Src-family kinases Lck and Fyn at their conserved negative regulatory carboxy-terminal tyrosine residues and upregulates kinase activity. Conversely, studies indicate that CD45 can also inhibit

tyrosine residues and upregulates kinase activity. Conversely, studies indicate that CD45 can also inhibit Lck and Fyn by dephosphorylating their positive regulatory autophosphorylation site. CD45 appears to be both a positive and a negative regulator that conducts signals depending on specific stimuli and cell type (1). Human leukocytes including lymphocytes, eosinophils, monocytes, basophils, and neutrophils express CD45, while erythrocytes and platelets are negative for CD45 expression (4).

CD45, write erythrocytes and platelets are negative for CD45 expression (4).

Background References 1. Huntington, N.D. and Tarlinton, D.M. (2004) *Immunol Lett* 94, 167-74.

2. Felberg, J. and Johnson, P. (2000) Biochem Biophys Res Commun 271, 292-8.

3. Kashio, N. et al. (1998) J Biol Chem 273, 33856-63.

4. Wang, Y. and Johnson, P. (2005) J Biol Chem 280, 14318-24.

Species Reactivity Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Applications Key FC-FP: Flow Cytometry (Fixed/Permeabilized)

Cross-Reactivity Key H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster

X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse

GP: Guinea Pig Rab: rabbit All: all species expected

Trademarks andCell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

Patents All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more

information.

Limited UsesExcept as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the

following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in

writing by a legally authorized representative of CST, are rejected and are of no force or effect.

1/1/24. 9:12 AM

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.